HARMONIC FUNCTIONS FOR SINGULAR QUADRANT WALKS

VIET HUNG HOANG, KILIAN RASCHEL, AND PIERRE TARRAGO

ABSTRACT. We consider discrete (time and space) random walks confined to the quarter plane, with jumps only in directions (i, j) with $i + j \ge 0$ and small negative jumps, i.e., $i, j \ge -1$. These walks are called singular, and were recently intensively studied from a combinatorial point of view. In this paper, we show how the compensation approach introduced in the 90ies by Adan, Wessels and Zijm may be applied to compute positive harmonic functions with Dirichlet boundary conditions. In particular, in case the random walks have a drift with positive coordinates, we derive an explicit formula for the escape probability, which is the probability to tend to infinity without reaching the boundary axes. These formulas typically involve famous recurrent sequences, such as the Fibonacci numbers. As a second step, we propose a probabilistic interpretation of the previously constructed harmonic functions and prove that they allow to compute all positive harmonic functions of these singular walks. To that purpose, we derive the asymptotics of the Green functions in all directions of the quarter plane and use Martin boundary theory.

1. INTRODUCTION

Espace probability for random walks in cones. Consider a multidimensional lattice random walk $\{S(n)\}_{n\geq 1}$, i.e., $S(n) = X(1) + \cdots + X(n)$ for all $n \geq 1$. Given a cone $C \subset \mathbb{R}^d$, $d \geq 1$, introduce the associated first exit time

$$\tau_x = \inf\{n \ge 0 : x + S(n) \notin C\} \le \infty.$$

If the drift of the increment distribution belongs to the cone, then (ignoring pathological behaviours) for x interior to C, the escape probability (also called survival probability)

(1)
$$\mathbb{P}(\tau_x = \infty) = \mathbb{P}(\forall n \ge 0, x + S(n) \in C)$$

is strictly positive, and defines a discrete harmonic function with Dirichlet boundary conditions.

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under the Grant Agreement No. 759702, from Centre Henri Lebesgue (programme ANR-11-LABX-0020-0), from the ANR DeRerumNatura (ANR-19-CE40-0018) and ANR CORTIPOM (ANR-21-CE40-0019).

Date: August 2, 2022.

²⁰²⁰ Mathematics Subject Classification. Primary 31C35, 60G50; Secondary 60J45, 60J50, 31C20, 11B39.

Key words and phrases. Discrete harmonic functions, singular random walks in the quarter plane, compensation approach, Green functions, Martin boundary.



FIGURE 1. Various step sets. From left to right: the simplest singular random walk; an arbitrary small step singular random walk; an example of singular walk with bigger jumps.

The question at the origin of the present work is the following: does this natural harmonic function admit an expression in closed form? The harmonicity property is equivalent to a recurrence relation, which in dimension 1 may be easily solved, at least in the bounded jump case. On the other hand, it is known that the behaviour of solutions to multivariate recurrences is much harder and vast [5, 6], a fortiori with boundary conditions depending on a cone; there is no hope, in general, to compute explicitly the escape probability (1).

A glimpse of our results (Part 1). We may now state the contributions of our paper, which consists of two parts. We will introduce a class of singular random walks in dimension 2 (see the next subsection for a precise definition, see also Figure 1), and look at the case of the cone C being the positive quarter plane. In Part 1, we will produce explicit expressions for the escape probability (1). To that purpose, we will use the compensation approach, as introduced in [2, 1, 3] by Adan, Wessels and Zijm. Before giving more details both on the random walks considered and on the techniques used, let us present an explicit example.

Consider in this paragraph the model $p_{-1,1} = p_{1,1} = p_{1,-1} = \frac{1}{3}$, as on Figure 1 (left). To that example, our main result will entail that the escape probability starting at x = (i, j) equals

(2)
$$\mathbb{P}(\tau_{(i,j)} = \infty) = 1 - \frac{1}{2^i} - \frac{1}{2^j} + \frac{1}{2^i 5^j} + \frac{1}{2^j 5^i} - \frac{1}{5^i 13^j} - \frac{1}{5^j 13^i} + \frac{1}{13^i 34^j} + \frac{1}{13^j 34^i} + \cdots$$

The integers appearing in the denominators in (2), namely $1, 2, 5, 13, 34, 89, 233, 610, \ldots$ are directly related to Fibonacci sequence (A000045), as the *n*-th term in the sequence is equal to F_{2n-1} . For i = j = 1, this result is derived in [24, Prop. 9], using a functional equation approach.

In this paper, we will prove similar formulas for other singular random walks; in particular, we will see how to define the sequence of denominators in general (providing a nice interplay between probabilistic and arithmetic properties of these singular random walks). As a second step, we will prove that similar expressions hold for infinitely many positive harmonic functions, of the form

(3)
$$h(i,j) = \sum_{n \ge 0} c_n \alpha_n^i \beta_n^j,$$

where the α_n, β_n, c_n are real constants (to be specified).

A glimpse of our results (Part 2). The main objective of our second part is to propose a probabilistic interpretation of the harmonic functions (3) constructed via the compensation approach. Our central result is to prove that the previous harmonic functions actually allow to construct all positive harmonic functions. In other words, the compensation approach yields an exhaustive description of positive harmonic functions for singular random walks. In concrete terms, this means that there is a correspondence between minimal positive harmonic functions and the yellow domain on the left display on Figure 4, exactly the same phenomenon as in the non-singular case [20]. This result is properly stated in Corollary 20.

Among all positive harmonic functions, there is a unique bounded positive harmonic function, which is the escape probability. It corresponds asymptotically by looking at the Green function

(4)
$$G(x,y) = \sum_{n \ge 1} \mathbb{P}(x+S(n)=y, \tau_x > n)$$

along the drift direction.

To prove these results, we use Martin boundary theory, and we believe that several intermediate results are of independent interest. The key idea is to compute the asymptotics of the Green function (4) as x is fixed and y goes to infinity in any direction of the cone. See Theorem 11 for the main statement. While such results have been recently derived in a close context (zero drift random walks [15], irreducible non-zero drift random walks [20]), there was no version in the literature applying to our context. The case of a boundary direction needs a particular attention, due to the interaction with the axes.

From a technical point of view, Part 2 is inspired by the work [20], [11] and [15].

Singular random walks in the quadrant. Throughout the paper, we assume that the distribution of the increments X(i) of the random walk have transition probabilities $p_{i,j}$ in \mathbb{Z}^2 such that (see Figure 1):

- (i) $\sum_{i,j} p_{i,j} = 1$ (normalization);
- (ii) $p_{i,j} = 0$ for all $i \leq -2$ or $j \leq -2$ (small negative jumps);
- (iii) $p_{-1,-1} = p_{-1,0} = p_{0,-1} = 0$ (singular walks);
- (iv) $p_{-1,1}p_{1,-1} \neq 0$;
- (v) There exists (i, j) with i + j > 0 such that $p_{i,j} > 0$ (non-degeneracy);
- (vi) The $p_{i,j}$ admit exponential moments in the following sense: the Laplace transform $\sum_{i,j} p_{i,j} e^{ix+jy}$ is finite in a neighborhood of any point of the curve (11) (moment assumption).

By definition, the kernel of the model is the bivariate polynomial

(5)
$$K(\alpha,\beta) = \alpha\beta \left(\sum_{i,j \ge -1} p_{i,j}\alpha^i\beta^j - 1\right).$$

Compensation approach. This technique has been developed in the probabilistic context of stationary distributions for random walks, see [2, 1, 3]. It does not aim directly at obtaining a solution for a generating function (as it is usual for quadrant walk problems), but rather tries to find a solution for its coefficients, in our case the escape probability $h(i, j) = \mathbb{P}(\tau_{(i,j)} = \infty)$ starting from (i, j).

As a discrete harmonic function, h(i, j) satisfies certain recursion relations (coming from harmonicity in our case), which differ depending on whether the state (i, j) lies on the boundary or not:

(6)
$$h(i,j) = \sum_{k,\ell \ge -1} p_{k,\ell} h(i+k,j+\ell), \quad \forall i,j \ge 1$$

(7)
$$h(i,0) = 0, \quad \forall i \ge 0,$$

(8)
$$h(0,j) = 0, \quad \forall j \ge 0.$$

The idea is then to express h(i, j) as a linear combination of products $\alpha^i \beta^j$, for pairs (α, β) such that the recursion relations (6) in the interior of the quarter plane hold. This is equivalent to choosing the parameters such that $K(\alpha, \beta) = 0$, with K as in (5). The products have to be chosen such that the recursion relations on the boundaries (7) and (8) are satisfied as well. As it turns out, this can be done by alternatingly compensating for the errors on the two boundaries, which eventually leads to an infinite series of product forms. The typical outcome of the compensation approach is an expression of the form (3). This clearly formally contains the series presented in (2).

As a side note, our work proposes a new example of applicability of the compensation approach, in relation with potential theory and discrete harmonic functions.

Related literature on explicit formulas for escape probabilities. In the quarter plane, the few known formulas concern non-singular walks with certain finite reflection groups (for which an algebraic version of the reflection principle applies); see Figure 2 for three examples. For instance, for the simple random walk (leftmost display on the figure), the escape probability equals [21, Cor. 8]

$$\mathbb{P}(\tau_{(i,j)} = \infty) = \left(1 - \left(\frac{p_{-1,0}}{p_{1,0}}\right)^i\right) \left(1 - \left(\frac{p_{0,-1}}{p_{0,1}}\right)^j\right),$$

which interestingly corresponds to a finite (four terms) sum in (3). Similar expressions hold for the other two models on Figure 2, with a sum involving six and eight terms, respectively.



FIGURE 2. Three non-singular step sets with finite reflection groups. The rightmost example is assumed to satisfy $p_{-1,1}p_{1,-1} = p_{-1,0}p_{1,0}$.

Still in the non-singular case, other approaches based on complex analysis techniques (boundary value problems) allow to compute the generating functions

$$\sum_{i,j \geqslant 1} h(i,j) x^i y^j$$

of harmonic functions in terms of certain conformal mappings, see [19] (zero drift case) and [22] (non-zero drift case).

In the continuous setting (Brownian motion in cones), there is a unified formula [18, Thm C] for the probability of escape from arbitrary cones $C \subset \mathbb{R}^d$, $d \ge 1$. More precisely, if the drift \vec{d} is interior to the cone, then

$$\mathbb{P}(\tau_x = \infty) = (2\pi)^{d/2} \exp\left(\left|x - \vec{d}\right|^2\right) p^C(1, x, \vec{d}),$$

where p^{C} denotes the heat kernel of the cone. The question of finding explicit expressions is then reduced to computing in closed-form the heat kernel, which is more classical. For example, if the cone is a Weyl chamber of type A, the determinantal Karlin-McGregor formula holds, and the survival probability is a finite sum of product forms, as in (3).

Applications and refinements of our results. We conclude the introduction by mentioning related open questions and potential applications of our results.

- One advantage of the compensation approach is that it is not based on generating functions. However, it would be interesting to understand how the compensation approach is related to Cohen and Boxma techniques [9, 8], which we developed in the paper [19] to study harmonic functions for non-singular random walks.
- In principle, the compensation approach could be applied as well to harmonic functions for reflected random walks in the quarter plane (with Neumann boundary conditions). It should also provide explicit expressions for *t*-harmonic functions (Dirichlet or Neumann boundary conditions).
- Formulas of the type of (3) hold for singular random walks (our main result) and for a few other (finite group) models, as those represented on Figure 2. This clearly suggests the question of determining the class of models for which harmonic functions may be expressed as in (3), via the pattern:

model \rightarrow curve \rightarrow sequence of numbers \rightarrow harmonic function (via Ansatz (3)).

- Is it possible to study more generally the Martin boundary and the Green functions asymptotics of singular random walks in arbitrary cones? What is the exact applicability of the compensation approach? For instance, can it be applied to harmonic functions of higher dimensional models?
- The compensation formula (15) that we find in the context of harmonic functions for singular random walks in the quarter plane is reminiscent of the alternating formula for the harmonic function of a space-time Brownian motion $(t, B_t)_{t \ge 0}$ conditioned to stay in the cone $\{(t, y), 0 < y < t\}$, see [10]. This suggests a relation between the singular walks we are considering and stochastic processes conditioned to stay in affine Weyl chambers.

Part 1. Constructing harmonic functions via the compensation approach

2. Properties of the Laplace transform and convergence of the series

This section aims at constructing harmonic functions as series taking the form of (3), by using the compensation approach. We first introduce a curve $\mathcal{K} \in \mathbb{R}^2$ such that $\alpha^i \beta^j$ is a solution of (6) for any $(\alpha, \beta) \in \mathcal{K}$, see Section 2.1. In Section 2.2, we then construct sequences $(\alpha_n, \beta_n)_{n \ge 0} \subset \mathcal{K}$ such that their series satisfies the boundary conditions (7)–(8). We then prove that these series converge and do not depend on the initial starting point of the sequence, see Section 2.3. Finally, in Section 2.4, we study a particular case, where we should renormalize the harmonic function to get a non-zero quantity.

2.1. Level sets of the Laplace transform. We first observe that the product form $\alpha^i \beta^j$ is solution to (6) if and only if

(9)
$$\alpha\beta = \sum_{k,\ell} p_{k,\ell} \alpha^{k+1} \beta^{\ell+1},$$

or equivalently, using the kernel notation (5), if and only if $K(\alpha, \beta) = 0$. We therefore introduce the algebraic curve

(10)
$$\mathcal{K} := \{ (\alpha, \beta) \in \mathbb{R}^2_{\geq 0} : K(\alpha, \beta) = 0 \}.$$

See Figure 3 for an example.

To take advantage of some convexity properties, we mainly investigate \mathcal{K} through an alternative exponential scaling as follows:

(11)
$$\mathcal{G} := \{ (x, y) \in \mathbb{R}^2 : K(e^x, e^y) = 0 \}.$$

See Figures 3 and 4 for examples of curves \mathcal{G} . Let us further denote

(12) $\mathcal{G}^+ := \{(x,y) \in \mathbb{R}^2 : K(e^x, e^y) < 0\}$ and $\mathcal{G}^- := \{(x,y) \in \mathbb{R}^2 : K(e^x, e^y) > 0\}.$

The following lemma presents some crucial properties of the curve \mathcal{G} .

Lemma 1. Under Assumptions (i)–(vi), we have the following assertions:

- (i) \mathcal{G}^+ is an unbounded convex domain and includes the ray $\{t(1,1): t < 0\}$;
- (ii) \mathcal{G} passes through (0,0) and the two tails of \mathcal{G} lie in the third quadrant \mathbb{R}^2_- ;
- (iii) \mathcal{G} admits a tangent at (0,0) satisfying the equation

$$\left(\sum_{k,\ell} kp_{k,\ell}\right) x + \left(\sum_{k,\ell} \ell p_{k,\ell}\right) y = 0.$$

Proof. We first prove (i). $K(e^x, e^y)$ can be factorized as

$$K(e^x, e^y) = e^{x+y}G(x, y),$$

where G(x, y) is defined as

(13)
$$G(x,y) := \sum_{k,\ell} p_{k,\ell} e^{kx+\ell y} - 1.$$

Since $\sum_{k,\ell} p_{k,\ell} e^{kx+\ell y}$ is a moment-generating function of a random variable in \mathbb{Z}^2 , it is convex in \mathbb{R}^2 . The domain \mathcal{G}^+ is thus convex. By letting x = y < 0, we then have:

$$G(x,x) = \sum_{k,\ell} p_{k,\ell} e^{(k+\ell)x} - 1 < \sum_{k,\ell} p_{k,\ell} - 1 = 0,$$

since $p_{k,\ell} = 0$ if $k + \ell < 0$. Thus, \mathcal{G}^+ is unbounded and includes the ray $\{t(1,1) : t < 0\}$.

We now prove (ii). Since G(0,0) = 0, then $(0,0) \in \mathcal{G}$. By letting x = 0 > y with |y| large enough, we have

$$G(0,y) = \sum_{k,\ell} p_{k,\ell} e^{\ell y} - 1 > p_{1,-1} e^{-y} > 0.$$

Similarly, for all y = 0 > x with |x| large enough, we have

$$G(x,0) = \sum_{k,\ell} p_{k,\ell} e^{kx} - 1 > p_{-1,1} e^{-x} > 0.$$

It implies that both tails of \mathcal{G} lie in \mathbb{R}^2_- .

Finally, (iii) is easily seen from the equation of the tangent at (0,0), which is

$$\partial_x K(e^0, e^0) x + \partial_y K(e^0, e^0) y = 0.$$

Although \mathcal{G}^+ is always convex, the bounded domain delimited by \mathcal{K} is not necessarily convex (see Figure 3). This is the main reason for us to use the curve \mathcal{G} rather than \mathcal{K} .

We now parametrize the curve \mathcal{G} by functions. Define $f : \mathbb{R}_{\leq 0} \to \mathbb{R}$ such that for all $x \leq 0$, $K(e^x, e^{f(x)}) = 0$ and $f(x) \geq x$. Similarly, we define $g : \mathbb{R}_{\leq 0} \to \mathbb{R}$ such that $K(e^{g(y)}, e^y) = 0$ and $g(y) \geq y$ for all $y \leq 0$. The following lemma describes key properties of f and g.

Lemma 2. Under Assumptions (i)–(vi), we have the following assertions:

(i) f and g are well defined, concave, and infinitely differentiable on \mathbb{R}_{-} ;



FIGURE 3. The model $p_{1,1} = 5/6$, $p_{1,-1} = p_{-1,1} = 1/12$: the interior of \mathcal{K} (left) is not convex, while the interior of \mathcal{G} (right) is convex.

- (ii) $\max_{x \leq 0} f(x) > f(0) = 0$ and $\arg \max_{x \leq 0} f(x)$ includes a unique point, denoted by x_0 . Further, f(x) is strictly increasing on $(-\infty, x_0)$, strictly decreasing on $(x_0, 0)$, and $\lim_{x \to -\infty} f(x) = -\infty$;
- (iii) $\max_{y \leq 0} g(y) > g(0) = 0$ and $\arg \max_{y \leq 0} g(y)$ includes a unique point, denoted by y_0 . Further, g(y) is strictly increasing on $(-\infty, y_0)$, strictly decreasing on $(y_0, 0)$, and $\lim_{y \to -\infty} g(y) = -\infty$.

Proof. We first prove Item (i). By the convexity of \mathcal{G}^+ , the slope of \mathcal{G} at (0,0), and the tails' position of \mathcal{G} (see Lemma 1), it is easily seen that f and g are well defined and concave on $\mathbb{R}_{\leq 0}$. Since any point $(x, y) \in \mathcal{G}$ is not the minimiser of the convex function G defined in (13), then $\partial_x G$ and $\partial_y G$ cannot vanish simultaneously for any $x, y \in \mathcal{G}$. By the construction of f and g, we then have

$$\partial_x G(x, f(x)) \neq 0$$
 and $\partial_y G(g(y), y) \neq 0$

for all $x, y \leq 0$. Thus,

$$f'(x) = -\frac{\partial_x G(x, f(x))}{\partial_y G(x, f(x))}$$
 and $g'(y) = -\frac{\partial_y G(g(y), y)}{\partial_x G(g(y), y)}$

are well defined for all x, y < 0, and so are all the higher orders derivatives of f and g.

We now prove Item (ii). Since

$$f'(0) = -\frac{\partial_x K}{\partial_y K}(e^0, e^0) = \frac{\sum_{k,\ell} k p_{k,\ell}}{\sum_{k,\ell} \ell p_{k,\ell}} < 0,$$

then $\max_{x \leq 0} f(x) > f(0) = 0$. If $\arg \max_{x \leq 0} f(x)$ includes at least two distinct points, then it also contains the segment between these points by the concavity of f. This implies that $K(\exp(x), \exp(\max_{x \leq 0} f(x)))$ is a constant function of x, which does not hold true. Hence, $\arg \max_{x \leq 0} f(x)$ includes a unique point x_0 . The strict monotonicity of f on (∞, x_0) and $(x_0, 0)$ then follows. By letting $x_1 < x_0$ such that $f(x_1) = 0$, we have for all $x < x_1$,

$$f(x) < \frac{f(x_0)}{x_0 - x_1} (x - x_1) \xrightarrow{x \to -\infty} -\infty.$$



FIGURE 4. Left: an example of curve \mathcal{G} (blue color) and the subpart (yellow) \mathcal{G}_0 corresponding to the Martin boundary. Right: an example of the compensation approach procedure.

Thus, $f(x) \to -\infty$ as $x \to -\infty$.

Item (iii) is proven similarly to (ii). The proof is then complete.

2.2. Construction of the product forms. Our main objective here is to prove Lemma 3 below, i.e., to construct sequences $\{(a_n, b_n)\}_{n \in \mathbb{Z}} \subset \mathcal{G}$ such that, setting

(14)
$$(\alpha_n, \beta_n) := \left(e^{a_n}, e^{b_n}\right) \in \mathcal{K},$$

the associated series

(15)
$$h(i,j) := \sum_{n \in \mathbb{Z}} \left(\alpha_n^i \beta_n^j - \alpha_{n+1}^i \beta_n^j \right)$$

is well defined, positive and satisfies (6)-(8).

To that purpose, we introduce a few useful notations. We first define

(16)
$$\mathcal{G}_0 := \{ (x, f(x)) : x \in (x_0, 0] \} \cup \{ (g(y), y) : y \in (y_0, 0] \} \subset \mathcal{G},$$

which corresponds to the yellow part on Figure 4. We first set

$$\widehat{f} := (f_{|(-\infty,x_0]})^{-1}$$
 and $\widetilde{f} := (f_{|(0,x_0]})^{-1}$.

Since f is concave, strictly increasing on $(-\infty, x_0]$, then \widehat{f} is a well-defined function, convex and strictly increasing on $(-\infty, f(x_0)]$. Similarly, the function $\widehat{g} := (g_{|(-\infty, y_0]})^{-1}$ is convex, strictly increasing on $(-\infty, g(y_0)]$.

We construct a sequence $\{(a_n, b_n)\}_{n \in \mathbb{Z}}$ contained in \mathcal{G} as follows:

$$(17) \qquad (a_0, b_0) \in \mathcal{G}_0,$$

(18)
$$(a_n, b_n) := \left(\widehat{f} \circ (\widehat{g} \circ \widehat{f})^{\circ (n-1)}(b_0), (\widehat{g} \circ \widehat{f})^{\circ (n)}(b_0)\right), \qquad n \ge 1,$$

(19)
$$(a_{-n}, b_{-n}) := \left((\widehat{f} \circ \widehat{g})^{\circ(n)}(a_0), \widehat{g} \circ (\widehat{f} \circ \widehat{g})^{\circ(n-1)}(a_0) \right), \qquad n \ge 1$$

Lemma 3. Let $\{(a_n, b_n)\}_{n \in \mathbb{Z}}$ be any sequence defined by (17), (18) and (19), and set $(\alpha_n, \beta_n) = (e^{a_n}, e^{b_n})$ as in (14). Then the series given by (15) converges absolutely for any $i, j \ge 1$ and satisfies Eq. (6)–(8). Furthermore, $h(i, j) \ge 0$ for any $i, j \ge 1$.

Proof. We start by proving the convergence of the series. To that aim, we construct an auxiliary sequence $(\hat{a}_n, \hat{b}_n)_{n \in \mathbb{Z}}$ as follows:

$$(\hat{a}_0, \hat{b}_0) = (a_0, b_0) \in \mathcal{G}_0,$$

$$(\hat{a}_n, \hat{b}_n) = (b_0 - nc_1 - (n-1)c_2, b_0 - nc_1 - nc_2), \qquad n \ge 1,$$

$$(\hat{a}_{-n}, \hat{b}_{-n}) = (a_0 - nc_1 - nc_2, a_0 - (n-1)c_1 - nc_2), \qquad n \ge 1,$$

where we have set $c_1 = f(x_0) - x_0 > 0$ and $c_2 = g(y_0) - y_0 > 0$, with x_0, y_0 as in Lemma 2. Let us also put, for any $n \in \mathbb{Z}$,

$$(\widehat{\alpha}_n, \widehat{\beta}_n) = (e^{\widehat{a}_n}, e^{\widehat{b}_n}).$$

Since \mathcal{G}^+ (see (12)) is convex and contains the ray $\{t(1,1): t < 0\}$, it should also contain the rays $\{(x_0, f(x_0)) + t(1,1): t < 0\}$ and $\{(g(y_0), y_0) + t(1,1): t < 0\}$. This implies that

(20)
$$\widehat{f}(y) \leqslant y - c_1 \quad \text{and} \quad \widehat{g}(x) \leqslant x - c_2,$$

for all $y \leq f(x_0)$ and $x \leq g(y_0)$. Thanks to the above inequalities and the monotonicity of \hat{g} , we first have

$$a_1 \leqslant \widehat{a}_1 \quad \text{and} \quad b_1 = \widehat{g}(a_1) \leqslant \widehat{g}(\widehat{a}_1) \leqslant \widehat{b}_1.$$

By induction argument, we then have

$$a_n = \widehat{f}(b_{n-1}) \leqslant \widehat{f}(\widehat{b}_{n-1}) \leqslant \widehat{a}_n$$
 and $b_n = \widehat{g}(a_{n-1}) \leqslant \widehat{g}(\widehat{a}_{n-1}) \leqslant \widehat{b}_n$

for any $n \ge 2$. Similarly, $a_{-n} \le \widehat{a}_{-n}$ and $b_{-n} \le \widehat{b}_{-n}$ for any $n \ge 1$. We now have

$$\sum_{n \in \mathbb{Z}} (\alpha_n^i \beta_n^j + \alpha_{n+1}^i \beta_n^j) \leqslant \sum_{n \in \mathbb{Z}} (\widehat{\alpha}_n^i \widehat{\beta}_n^j + \widehat{\alpha}_{n+1}^i \widehat{\beta}_n^j) = \widehat{\alpha}_0^i \widehat{\beta}_{-1}^j + \widehat{\alpha}_0^i \widehat{\beta}_0^j + \widehat{\alpha}_1^i \widehat{\beta}_0^j + \sum_{n \ge 1} (e^{-jc_2} + e^{jc_1}) \alpha_0^{i+j} \left(e^{-(c_1 + c_2)(i+j)} \right)^n + \sum_{n \ge 1} (e^{ic_2} + e^{-ic_1}) \beta_0^{i+j} \left(e^{-(c_1 + c_2)(i+j)} \right)^n.$$

The last sums are finite for any $i, j \ge 1$. Hence h(i, j) converges absolutely for any $i, j \ge 1$.

We now prove that the series h(i, j) satisfies Eq. (6)–(8). Firstly, all the terms $\alpha_n^i \beta_n^j$ and $\alpha_{n+1}^i \beta_n^j$ satisfy Eq. (6) (since (α_n, β_n) and $(\alpha_{n+1}, \beta_n) \in \mathcal{K}$), then h(i, j) should also satisfy Eq. (6). Secondly, since

$$h(i,0) = \sum_{n \in \mathbb{Z}} (-\alpha_n^i + \alpha_n^i) = 0 \quad \text{and} \quad h(0,j) = \sum_{n \in \mathbb{Z}} (\beta_n^j - \beta_n^j) = 0,$$

then h(i, j) satisfies Eq. (7) and (8).

We now prove that $h(i, j) \ge 0$ for all values of $i, j \ge 1$, by rewriting

$$h(i,j) = \alpha_0^i \beta_0^j \left(1 - \left(\frac{\beta_{-1}}{\beta_0}\right)^j - \left(\frac{\alpha_1}{\alpha_0}\right)^i \right) + \sum_{n \ge 1} \alpha_{-n}^i \left(\beta_{-n}^j - \beta_{-(n+1)}^j\right) + \sum_{n \ge 1} \left(\alpha_n^i - \alpha_{n+1}^i\right) \beta_n^j.$$

The first term is positive for any $i, j \ge 1$ such that i+j is large enough, whereas the second and third terms are positive since $\alpha_{-n} > \beta_{-(n+1)} > \alpha_{-(n+1)}$ and $\beta_n > \alpha_{n+1} > \beta_{n+1}$ for any $n \ge 0$. Hence, there exists $n_0 > 0$ such that h(i, j) > 0 for all $i, j \ge 1$ and $i+j \ge n_0$. Thus,

$$h(i,j) = \sum_{k,\ell} p_{k,\ell} h(i+k,j+\ell) \ge 0$$

for any $i, j \ge 1$ and $i + j = n_0 - 1$. By induction argument, we have $h(i, j) \ge 0$ for all $i, j \ge 1$. The proof is then complete.

The sign of harmonic functions h(i, j) can be asserted more specifically.

Remark 4. If the walk has jumps inside the positive quadrant, i.e., there exist $k, \ell \ge 0$ such that $k + \ell > 0$ and $p_{k,\ell} > 0$, then h(i,j) > 0 for all $i, j \ge 1$. On the other hand, if the walk does not have any jumps inside the positive quadrant, i.e., $p_{k,\ell} = 0$ for all $k, \ell \ge 0$, then h(i,j) > 0 for $i, j \ge 1$ and i + j large enough, but h(i,j) can vanish for some small i, j.

2.3. Dependency on the starting point. In Lemma 3, we assumed that the starting point (a_0, b_0) belongs to \mathcal{G}_0 . In this subsection, we prove that we do not obtain more harmonic functions with starting points in $\mathcal{G} \setminus \mathcal{G}_0$.

We construct a sequence $\{(a_n, b_n)\}_{n \in \mathbb{Z}}$ as follows:

 $(21) (a_0, b_0) \in \mathcal{G},$

(22) $a_n \neq a_{n-1}$ s.t. $(a_n, b_{n-1}) \in \mathcal{G}, \quad b_n \neq b_{n-1}$ s.t. $(a_n, b_n) \in \mathcal{G},$

(23) $b_{-n} \neq b_{-(n-1)}$ s.t. $(a_{-(n-1)}, b_n) \in \mathcal{G}, \quad a_{-n} \neq a_{-(n-1)}$ s.t. $(a_{-n}, b_{-n}) \in \mathcal{G}, \quad n \ge 1.$

Let us define

$$\mathcal{O}(\mathcal{G}_0) := \{\{(a_n, b_n)\}_{n \in \mathbb{Z}} \text{ satisfying } (21) - (23) \text{ s.t. } (a_0, b_0) \in \mathcal{G}_0\}.$$

In other words, $\mathcal{O}(\mathcal{G}_0)$ is the set of orbits formed by the sequences $\{(a_n, b_n)\}_{n \in \mathbb{Z}}$ satisfying (21)–(23) and initiated from a point in \mathcal{G}_0 .

We have the following lemma.

Lemma 5. Any sequence $\{(a_n, b_n)\}_{n \in \mathbb{Z}}$ satisfying (21)–(23) belongs to the set $\mathcal{O}(\mathcal{G}_0)$. As a result, the series h(i, j) defined in (15) does not depend on the initial value (α_0, β_0) , but only depends on the orbits in $\mathcal{O}(\mathcal{G}_0)$.

Proof. Assume that $\{(a_n, b_n)\}_{n \in \mathbb{Z}}$ satisfies (21)–(23) with $(a_0, b_0) \notin \mathcal{G}_0$. It is sufficient to show that there always exists $n_0 \in \mathbb{Z}$ such that $(a_{n_0}, b_{n_0}) \in \mathcal{G}_0$ or $(a_{n_0}, b_{n_0-1}) \in \mathcal{G}_0$.

If $0 \leq a_0 < g(y_0)$, then $(a_0, b_{-1}) = (a_0, (g|_{[y_0,0]})^{-1}(a_0)) \in \mathcal{G}_0$. Similarly, if $0 \leq b_0 < f(x_0)$, then $(a_1, b_0) = ((f|_{[x_0,0]})^{-1}(b_0)) \in \mathcal{G}_0$.

If $x_0 < a_0 < 0$, then $(a_0, b_{-1}) = (a_0, f(a_0)) \in \mathcal{G}_0$. If $y_0 < b_0 < 0$, $(a_1, b_0) = (g(b_0), b_0) \in \mathcal{G}_0$.

If $a_0 < x_0$ and $b_0 < y_0$, without loss of generality, we will further assume that $b_0 = f(a_0)$. Since

 $f(x) \ge x + c_1$ and $g(y) \ge y + c_2$,

for all $x \leq x_0$ and $y \leq y_0$, then there exists $n_0 \geq 1$ such that $a_{n_0} = (g \circ f)^{\circ(n)}(a_0) > x_0$ or $b_{n_0} = f \circ (g \circ f)^{\circ(n)}(a_0) > y_0$. Hence, a_{n_0} or b_{n_0} falls into the preceding cases. The rest of the proof then follows trivially.

2.4. Behavior on the boundary. We will show that the harmonic functions defined in (15), considered as a function of $(a_0, b_0) \in \mathcal{G}_0$, will converge (after normalization) to a non-trivial positive harmonic function as (a_0, b_0) tends to the boundary of \mathcal{G}_0 , which consists of the points $(x_0, f(x_0))$ and $(g(y_0), y_0)$, with x_0 and y_0 introduced in Lemma 2.

For that matter, let us re-introduce the sequence $\{(a_n, b_n)\}_{n \in \mathbb{Z}}$ in (17)–(19) as functions of one variable y:

$$\begin{split} b_{0}(y) &= y \in (y_{0}, 0), \quad a_{0}(y) = g(y), \quad b_{-1}(y) = \widehat{g} \circ g(y), \\ (a_{n}(y), b_{n}(y)) &= \left(\widehat{f} \circ (\widehat{g} \circ \widehat{f})^{\circ (n-1)}(y), (\widehat{g} \circ \widehat{f})^{\circ (n)}(y)\right), \qquad n \ge 1, \\ (a_{-n}(y), b_{-(n+1)}(y)) &= \left(\widehat{f} \circ (\widehat{g} \circ \widehat{f})^{\circ (n-1)} \circ b_{-1}(y), (\widehat{g} \circ \widehat{f})^{\circ (n)} \circ b_{-1}(y)\right), \qquad n \ge 1. \end{split}$$

We further denote

$$(\alpha_n(y), \beta_n(y)) = \left(e^{a_n(y)}, e^{b_n(y)}\right), \quad n \in \mathbb{Z},$$
$$h_y(i, j) = \sum_{n \ge 0} \alpha_n(y)^i \beta_n(y)^j - \alpha_{n+1}(y)^i \beta_n(y)^j$$

for all $i, j \ge 0, y \in (y_0, 0)$. Following the above construction, we remark that

$$(a_{-n}(y), b_{-(n+1)}(y)) = (a_n \circ b_{-1}(y), b_n \circ b_{-1}(y))$$

for any $n \ge 1$. Then, putting for any $i, j \ge 0$ and $y \in (y_0, 0)$

$$t_{i,j}(y) = \sum_{n \ge 0} \alpha_n(y)^i \beta_n(y)^j - \alpha_{n+1}(y)^i \beta_n(y)^j,$$

one has

$$h_y(i,j) = t_{i,j}(y) - t_{i,j}(b_{-1}(y)).$$

The following proposition shows the convergence of harmonic functions $h_y(i,j)$ as $y \to y_0^+$.

Proposition 6. For any $i, j \ge 0$,

$$\lim_{y \to y_0^+} h_y(i,j) = 0$$

Moreover, for any $i, j \ge 0$, $t_{i,j}(y)$ is differentiable on $[y_0, 0)$ and

$$\lim_{y \to y_0^+} \frac{h_y(i,j)}{y - y_0} = 2t'_{i,j}(y_0).$$

Furthermore, $t'_{i,j}(y_0) \ge 0$ for all $i, j \ge 1$.

The remainder of this section is devoted to proving Proposition 6. We first study the derivatives of some important functions.

Lemma 7. We have:

- (i) $\lim_{y\to y_0^+} b_{-1}(y) = y_0$ and $b'_{-1}(y_0) = -1$;
- (ii) $\widehat{f'}$ is strictly positive and strictly increasing on $(-\infty, f(x_0))$. Furthermore,

$$\lim_{y \to -\infty} \widehat{f}'(y) = 1;$$

(iii) \hat{g}' is strictly positive and strictly increasing on $(-\infty, g(y_0))$. Furthermore,

$$\lim_{x \to -\infty} \widehat{g}'(x) = 1.$$

Proof. We first prove Item (i). Since g(y) is strictly concave on $(-\infty, 0)$ and admits its maximum value at y_0 , then $g'(y_0) = 0$ and

$$g(y) = g(y_0) + \frac{g''(y_0)}{2}(y - y_0)^2 + o((y - y_0)^2)$$

as $y \to y_0$, where $g''(y_0) < 0$. Hence,

$$\lim_{y \to y_0^+} \left(\frac{b_{-1}(y) - y_0}{b_0(y) - y_0} \right)^2 = \lim_{y \to y_0^+} \frac{g(b_{-1}(y)) - g(y_0)}{g(b_0(y)) - g(y_0)} = 1.$$

This implies $b_{-1}(y) \to y_0$ as $b_0(y) = y \to y_0^+$. Since $b_{-1}(y) < y_0 < b_0(y)$ for all $y \in (y_0, 0)$, then $b'_{-1}(y_0) = -1$.

We move to the proof of Item (ii) (Item (iii) would be proven similarly). Since \hat{f} is strictly convex and strictly increasing on $(-\infty, f(x_0))$, then \hat{f}' is strictly increasing and positive on $(-\infty, f(x_0))$. Putting

$$\alpha(\beta) = e^{f(\log \beta)}, \quad \beta \ge 0,$$

we first study the behavior of $\alpha'(\beta)$ as $\beta \to 0$. Since $\alpha(\beta)$ satisfies $K(\alpha(\beta), \beta) = 0$, one may differentiate once and twice the equation and evaluate them at $\beta = 0$ as follows:

$$\partial_{\alpha}K(0,0)\alpha'(0) + \partial_{\beta}K(0,0) = 0,$$

$$\partial_{\alpha\alpha}K(0,0)\alpha'(0)^{2} + 2\partial_{\alpha\beta}K(0,0)\alpha'(0) + \partial_{\beta\beta}K(0,0) + \partial_{\alpha}K(0,0)\alpha''(0) = 0.$$

While the value of $\alpha'(0)$ cannot be deduced from the first equation, $\alpha'(0)$ can be solved explicitly from the second:

$$\alpha'(0) = \frac{1 - p_{0,0} \pm \sqrt{(1 - p_{0,0})^2 - 4p_{1,-1}p_{-1,1}}}{2p_{1,-1}} \neq 0, \infty.$$

These solutions correspond to the behavior of the two branches $(e^{\hat{f}(y)}, e^y)$ and $(e^{g(y)}, e^y)$ as $y \to -\infty$. This implies that

$$\widehat{f}'(y) = \alpha'(e^y) \xrightarrow{e^y} \xrightarrow{y \to -\infty} \alpha'(0) \frac{1}{\alpha'(0)} = 1.$$

The following lemma shows the convergence of $t'_{i,j}(y)$.

Lemma 8. For any $i, j \ge 1$ and $y \in [y_0, 0)$, the series

$$t'_{i,j}(y) = \sum_{n \ge 0} \left(\alpha_n(y)^i \beta_n(y)^j - \alpha_{n+1}(y)^i \beta_n(y)^j \right)'$$

converges absolutely.

Proof. By Lemma 7, the derivatives of $\alpha_n(y)$ and $\beta_n(y)$ are well defined on $[y_0, 0)$ for all $n \ge 0$. Since $\hat{f}(y), \hat{g}(x) \downarrow 1$ as $x, y \to -\infty$, then

$$a'_{n}(y) = \hat{f} \circ b_{n-1}(y) \prod_{k=1}^{n-1} \hat{g}' \circ a_{k}(y) \cdot \hat{f}' \circ b_{k-1}(y) = o(r^{n}),$$

$$b'_{n}(y) = \prod_{k=1}^{n} \hat{g}' \circ a_{k}(y) \cdot \hat{f}' \circ b_{k-1}(y) = o(r^{n}),$$

as $n \to \infty$, for any r > 1. Let us now fix $i, j \ge 1$. Recall from the proof of Lemma 3 that

$$\alpha_n(y)^i \beta_n(y)^j \leqslant y^{i+j} e^{ic_2} \left(e^{-(i+j)(c_1+c_2)} \right)^n = O\left(e^{-(i+j)(c_1+c_2)} \right)^n,$$

as $n \to \infty$, where c_1 and c_2 are positive constants. Hence,

$$\left|\left(\alpha_n(y)^i\beta_n(y)^j\right)'\right| = \left|\left(ia'_n(y) + jb'_n(y)\right)\alpha_n(y)^i\beta_n(y)^j\right| \le o(r^n),$$

as $n \to \infty$, for any $y \in [y_0, 0)$ and r > 0. Similarly, we also have

$$\left|\left(\alpha_{n+1}(y)^{i}\beta_{n}(y)^{j}\right)'\right| \leqslant o(r^{n}),$$

as $n \to \infty$, for any $y \in [y_0, 0)$ and r > 0. Thus, the series

$$t'_{i,j}(y) = \sum_{n \ge 0} \left(ia'_n(y) + jb'_n(y) \right) \alpha_n(y)^i \beta_n(y)^j - \left(ia'_{n+1}(y) + jb'_n(y) \right) \alpha_{n+1}(y)^i \beta_n(y)^j$$

converges absolutely for any $i, j \ge 1$.

We now prove Proposition 6.

Proof of Proposition 6. Since $b_{-1}(y) \to y_0$ as $y \to y_0^+$ by Lemma 7(i), then

$$h_y(i,j) = t_{i,j}(y) - t_{i,j}(b_{-1}(y)) \to 0$$

as $y \to y_0^+$. By L'Hôpital's rule, we now deduce

$$\frac{h_y(i,j)}{y-y_0} \xrightarrow{y \to y_0^+} t'_{i,j}(y_0) - t'_{i,j}(b_{-1}(y_0))b'_{-1}(y_0) = 2t'_{i,j}(y_0).$$

Since $h_y(i, j)$ is harmonic and non-negative for all $y \in (y_0, 0)$, this guarantees that $t'_{i,j}(y_0)$ is harmonic and non-negative for all $i, j \ge 1$.

We now prove that $t'_{i,j}(y_0) > 0$ for all $i, j \ge 1$ and i + j large enough. Since for any $n \ge 0$ one has $\alpha_n(y_0) > \alpha_{n+1}(y_0)$, then

(24)
$$jb'_{n}(y_{0})\alpha_{n}(y_{0})^{i}\beta_{n}(y_{0})^{j} - jb'_{n}(y_{0})\alpha_{n+1}(y_{0})^{i}\beta_{n}(y_{0})^{j} > 0,$$

for any $n \ge 0$ and $i, j \ge 1$. We then show that

(25)
$$ia'_{n}(y_{0})\alpha_{n}(y_{0})^{i}\beta_{n}(y_{0})^{j} - ia'_{n+1}(y_{0})\alpha_{n+1}(y_{0})^{i}\beta_{n}(y_{0})^{j} > 0,$$

or equivalently,

$$\alpha_n(y_0)^i > \alpha_{n+1}(y_0)^i \left(\widehat{f}' \circ b_n(y_0)\right) \cdot \left(\widehat{g}' \circ a_n(y_0)\right),$$

for any $n \ge 0$, $j \ge 1$ and *i* large enough. Indeed, recall that

$$\frac{\alpha_n(y_0)}{\alpha_{n+1}(y_0)} = e^{a_n(y_0) - a_{n+1}(y_0)} \ge e^{f(x_0) - x_0 + g(y_0) - y_0} > 0,$$

for any $n \ge 0$, and $(\hat{f}' \circ b_n(y_0)) \cdot (\hat{g}' \circ a_n(y_0)) \downarrow 1$ as $n \to \infty$. Hence, for any $n \ge 0$ and i large enough,

$$\left(\frac{\alpha_n(y_0)}{\alpha_{n+1}(y_0)}\right)^i > \left(\widehat{f}' \circ b_n(y_0)\right) \cdot \left(\widehat{g}' \circ a_n(y_0)\right),$$

which is equivalent to Inequality (25). Both (24) and (25) imply that there exists $n_0 > 1$ such that $t'_{i,j}(y_0) > 0$ for any $j \ge 1$ and $i \ge n_0$. The harmonicity then implies that $t'_{i,j}(y_0) > 0$ for any $i, j \ge 1$ and $i + j \ge n_0 + 1$.

We want to point out that Remark 4 also holds true for the harmonic function $t_{i,j}(y_0)$, that is, $t_{i,j}(y_0) > 0$ for all $i, j \ge 1$ if there exist $k, \ell \ge 0$ such that $k + \ell \ge 1$ and $p_{k,\ell} > 0$.

3. Explicit expression for walks with small steps

This section aims at giving an explicit expression for all terms appearing in the harmonic functions (15), in the case where the random walks only have small jumps, that is, the positive transition probabilities can only be in the set $\{p_{-1,1}, p_{1,-1}, p_{1,0}, p_{0,1}, p_{1,1}\}$, see Figure 1.

The following lemma presents a uniformization of the zero set of $K(\alpha, \beta)$.

Lemma 9. One has

$$\{(\alpha,\beta)\in\mathbb{C}^2:K(\alpha,\beta)=0\}=\{(\alpha(s),\beta(s)):s\in\mathbb{C}\},\$$

where

$$\alpha(s)^{-1} := \frac{\sqrt{b^2 - ac}}{2a} \left(s + \frac{1}{s} \right) + \frac{b}{a} \quad and \quad \beta(s)^{-1} := \frac{\sqrt{\widehat{b^2} - a\widehat{c}}}{2a} \left(\rho s + \frac{1}{\rho s} \right) + \frac{\widehat{b}}{a},$$

and $a := 1 - 4p_{-1,1}p_{1,-1}$, $\rho := \sqrt{\frac{1+\sqrt{a}}{1-\sqrt{a}}}$, and finally

$$b := p_{0,1} + 2p_{-1,1}p_{1,0}, \qquad c := p_{0,1}^2 - 4p_{-1,1}p_{1,1},$$

$$\hat{b} := p_{1,0} + 2p_{1,-1}p_{0,1}, \qquad \hat{c} := p_{1,0}^2 - 4p_{1,-1}p_{1,1}.$$

We further have the involutions $\alpha(s) = \alpha(1/s)$ and $\beta(s) = \beta(1/(\rho^2 s))$.

Proof. We will find a rational uniformization of the algebraic curve

(26)
$$\alpha^2 \beta^2 K\left(\frac{1}{\alpha}, \frac{1}{\beta}\right) = 0.$$

We first rewrite

$$\alpha^{2}\beta^{2}K\left(\frac{1}{\alpha},\frac{1}{\beta}\right) = p_{1,1} + p_{1,0}\beta + p_{0,1}\alpha + p_{1,-1}\beta^{2} + p_{-1,1}\alpha^{2} - \alpha\beta$$
$$= (p_{1,-1})\beta^{2} + (-\alpha + p_{1,0})\beta + (p_{-1,1}\alpha^{2} + p_{0,1}\alpha + p_{1,1}).$$

The above polynomial admits the discriminant

$$\delta(\alpha) := (-\alpha + p_{1,0})^2 - 4p_{1,-1} \left(p_{-1,1}\alpha^2 + p_{0,1}\alpha + p_{1,1} \right)$$
$$= a\alpha^2 - 2b\alpha + c = \frac{b^2 - ac}{4a} \left[\left(2\frac{a\alpha - b}{\sqrt{b^2 - ac}} \right)^2 - 4 \right],$$

where a, b, c are defined in Lemma 9, and

$$b^{2} - ac = 4p_{1,-1} \left((1 - 4p_{-1,1}p_{1,-1})p_{1,1} + p_{1,0}^{2}p_{-1,1} + p_{0,1}^{2}p_{1,-1} + p_{0,1}p_{1,0} \right) > 0.$$

By setting

$$2\frac{a\alpha-b}{\sqrt{b^2-ac}} = s + \frac{1}{s}, \quad \text{that is,} \quad \alpha = \frac{\sqrt{b^2-ac}}{2a}\left(s + \frac{1}{s}\right) + \frac{b}{a}, \quad s \in \mathbb{C},$$

one can easily find β satisfying Eq. (26) as follows:

$$\beta = \frac{\alpha(s) - p_{0,1} + \sqrt{\delta \circ \alpha(s)}}{2p_{1,-1}} = \frac{\sqrt{\widehat{b}^2 - a\widehat{c}}}{2a} \left(\rho s + \frac{1}{\rho s}\right) + \frac{\widehat{b}}{a},$$

where \hat{b}, \hat{c}, ρ are defined in Lemma 9. The proof is then complete.

By the above lemma, one can describe the curve \mathcal{K} in (10) as

$$\mathcal{K} = \{ (\alpha(s), \beta(s)) : s > 0 \}.$$

We remark that at s = 1, the discriminant $\delta \circ \alpha(1) = 0$, then $\beta(1)$ is the double root of Eq. (26) as an equation of β . Similarly, at $s = 1/\rho$, $\alpha(1/\rho)$ is a double root of Eq. (26) as an equation of α . Thus, defining \mathcal{K}_0 as the analogue of \mathcal{G}_0 (see (16)) before the exponential change of variable, one easily deduces the description

$$\mathcal{K}_0 = \{ (\alpha(s), \beta(s)) : s \in (1/\rho, 1) \}$$

Proposition 10. Any sequence $\{(\alpha_n, \beta_n)\}_{n \in \mathbb{Z}}$ satisfying (21)–(23) and (14), with $(\alpha_0, \beta_0) \in \mathcal{K}_0$, can be described explicitly as

$$\{(\alpha_n,\beta_n)\}_{n\in\mathbb{Z}} = \left\{\left(\alpha(\rho^{2n}s),\beta(\rho^{2n}s)\right)\right\}_{n\in\mathbb{Z}},\$$

with $s \in (1/\rho, 1)$. As a consequence, the harmonic function formed by $(\alpha_n, \beta_n)_{n \in \mathbb{Z}}$ in (15) becomes

$$h(i,j) = \sum_{k \in \mathbb{Z}} \alpha(\rho^{2n}s)^i \beta(\rho^{2n}s)^j - \alpha(\rho^{2n+2}s)^i \beta(\rho^{2n}s)^j, \qquad i,j \ge 1.$$

16

Proof. We will prove the proposition by induction argument. Since $(\alpha_0, \beta_0) \in \mathcal{K}_0$, then

$$(\alpha_0, \beta_0) = (\alpha(s), \beta(s)),$$

with $s \in (1/\rho, 1)$. Now assume that $(\alpha_n, \beta_n) = (\alpha(\rho^{2n}s), \beta(\rho^{2n}s))$ with $n \in \mathbb{Z}$. Since $\beta_n = \beta(\rho^{2n}s) = \beta(1/(\rho^{2n}s))$ by the involution and (α_{n+1}, β_n) is a root of $K(\alpha, \beta)$, then

$$\alpha_{n+1} = \alpha \left(\frac{1}{\rho^{2n+2}s}\right) = \alpha(\rho^{2n+2}s).$$

Since $(\alpha_{n+1}, \beta_{n+1})$ is also a root of $K(\alpha, \beta)$, then $\beta_{n+1} = \beta(\rho^{2n+2}s)$. Similarly, the involutions of Lemma 9 also imply that $\alpha_{n-1} = \alpha(\rho^{2n-2}s)$ and $\beta_{n-1} = \beta(\rho^{2n-2}s)$. The proof is then complete.

A concrete example. We now return to our simplest walk with the transition probabilities $p_{1,1} = p_{1,-1} = p_{-1,1} = 1/3$ and provide a full proof of the formula (2). The zero set of the associated kernel

$$K(\alpha, \beta) = \frac{1}{3}(x^2y^2 + x^2 + y^2) - xy$$

admits the uniformization presented in Lemma 9, with

$$\alpha(s) = \frac{\sqrt{5}}{s+1/s}, \quad \beta(s) = \frac{\sqrt{5}}{\rho s+1/(\rho s)} \text{ and } \rho = \frac{3+\sqrt{5}}{2}.$$

The sequence $\{(\alpha_n, \beta_n)\}_{n \in \mathbb{Z}}$ in Proposition 10 can be expressed as functions of $s \in (1/\rho, 1)$:

$$(\alpha_n(s), \beta_n(s)) = \left(\frac{\sqrt{5}}{\rho^{2n}s + 1/(\rho^{2n}s)}, \frac{\sqrt{5}}{\rho^{2n+1}s + 1/(\rho^{2n+1}s)}\right), \qquad n \in \mathbb{Z}$$

Consequently, this leads to harmonic functions depending on s, denoted by

$$h_s(i,j) := \sum_{n \in \mathbb{Z}} \alpha_n(s)^i \beta_n(s)^j - \alpha_{n+1}(s)^i \beta_n(s)^j.$$

Let us look at two specific examples. Firstly, the survival probability (2) is the harmonic function associated to the initial point

$$\left(\alpha_0\left(\frac{\sqrt{5}-1}{2}\right),\beta_0\left(\frac{\sqrt{5}-1}{2}\right)\right) = (1,1) \in \mathcal{K}_0,$$

and can be expressed explicitly as

$$\mathbb{P}(\tau_{(i,j)} = \infty) = h_{\frac{\sqrt{5}-1}{2}}(i,j) = \dots - \frac{1}{5^i 13^j} + \frac{1}{5^i 2^j} - \frac{1}{1^i 2^j} + \frac{1}{1^i 1^j} - \frac{1}{2^i 1^j} + \frac{1}{2^i 5^j} - \frac{1}{13^i 5^j} + \dots$$

In this formula, the sequence

$$\left\{\frac{\rho^n s + \frac{1}{\rho^n s}}{\sqrt{5}}\right\}_{n \ge 0} = \{1, 1, 2, 5, 13, 34, 89, 233, 610, 1597, \ldots\}$$

is a bisection of Fibonacci numbers.

We now give the expression for the normalized positive harmonic function constructed in Proposition 6. Recall that such a function is obtained by differentiating the harmonic function $h_s(i, j)$ with respect to the variable $\log \beta_0(s)$ and evaluating as $b_0 \to \log \beta_0(1)$. (The logarithms appear in our computation because Proposition 6 was stated under $(\log \alpha, \log \beta)$ -coordinates instead of (α, β) -coordinates.) We have:

$$\frac{h_s(i,j)}{\log \beta_0(s) - \log \beta_0(1)} \xrightarrow{s \to 1^-} \left. \frac{dh_s(i,j)}{ds} \frac{ds}{d (\log \beta_0(s))} \right|_{s=1},$$

where

$$\begin{aligned} \frac{ds}{d(\log \beta_0(s))} \bigg|_{s=1} &= \frac{\beta_0(1)}{\beta'_0(1)} = -\frac{3}{\sqrt{5}}, \\ \frac{dh_s(i,j)}{ds} \bigg|_{s=1} \\ &= \sum_{n \in \mathbb{Z}} i\left(\alpha'_n(1)\alpha_n(1)^{i-1} - \alpha'_{n+1}(1)\alpha_{n+1}(1)^{i-1}\right)\beta_n(1)^j + j\left(\alpha_n(1)^i - \alpha_{n+1}(1)^i\right)\beta'_n(1)\beta_n(1)^{j-1}, \\ (\alpha_n(1), \beta_n(1)) &= \left(\frac{\sqrt{5}}{\rho^{2n} + \frac{1}{\rho^{2n}}}, \frac{\sqrt{5}}{\rho^{2n+1} + \frac{1}{\rho^{2n+1}}}\right), \\ (\alpha'_n(1), \beta'_n(1)) &= \left(-\frac{\sqrt{5}\left(\rho^{2n} - \frac{1}{\rho^{2n}}\right)}{\left(\rho^{2n} + \frac{1}{\rho^{2n}}\right)^2}, -\frac{\sqrt{5}\left(\rho^{2n+1} - \frac{1}{\rho^{2n+1}}\right)}{\left(\rho^{2n+1} + \frac{1}{\rho^{2n+1}}\right)^2}\right), \quad n \in \mathbb{Z}. \end{aligned}$$

In these formulas, we want to point out that the sequence

$$\left\{\rho^n + \frac{1}{\rho^n}\right\}_{n \ge 0} = \{2, 3, 7, 18, 47, 123, 322, 843, 2207, \ldots\} = \{L_{2n}\}_{n \ge 0}$$

is a bisection of Lucas numbers $\{L_n\}_{n \ge 0}$:

$$L_0 = 2, \quad L_1 = 1, \quad L_n = L_{n-1} + L_{n-2}, \quad n \ge 2,$$

see A005248. Moreover, the sequence

$$\{u_n\}_{n\geq 0} = \left\{\sqrt{5}\left(\rho^n - \frac{1}{\rho^n}\right)\right\}_{n\geq 0} = \{0, 5, 15, 40, 105, 275, 720, 1885, 4935, \ldots\}$$

satisfies the recurrence relation (see A201157):

$$u_0 = 0, \quad u_1 = 5, \quad u_n = 3u_{n-1} - u_{n-2}, \quad n \ge 2.$$

Part 2. Green functions and Martin boundary for singular random walks in the quadrant

In this part, we describe the Martin boundary inside the cone and prove that it coincides with the set of harmonic functions constructed in Part 1.

4. Presentation of the results

4.1. A brief account on Green functions and Martin boundary theory. Let us first recall the definition of the Green function of the random walk inside the cone $\mathbb{Z}_{>0}^2$, which plays an important role in the construction of the Martin boundary. For $x, y \in \mathbb{Z}_{>0}^2$, set

$$G(x,y) = \sum_{n \geqslant 1} \mathbb{P}(x + S(n) = y, \tau_x > n) \quad \text{and} \quad \widetilde{G}(x,y) = \sum_{n \geqslant 1} \mathbb{P}(x + S(n) = y),$$

where $\tau_x = \inf\{n \ge 0 : x + S(n) \notin \mathbb{Z}_{>0}^2\} \le \infty$, and set $x_0 = (1, 1)$. The Martin kernel associated to the reference point x_0 is then defined as

(27)
$$K_M(x,y) = \frac{G(x,y)}{G(x_0,y)}.$$

The Martin boundary $\partial_M^{\mathbb{Z}_{\geq 0}^2} S$ of the random walk S killed outside of $\mathbb{Z}_{\geq 0}^2$ is the boundary in the topological space $\{f : \mathbb{Z}_{\geq 0}^2 \to \mathbb{R}\}$ (with the topology of point-wise convergence) of the set of maps $\{K_M(\cdot, y) : y \in \mathbb{Z}_{\geq 0}^2\}$. The space $\partial_M^{\mathbb{Z}_{\geq 0}^2} S$ is a compact measurable subspace of the set of real functions on $\mathbb{Z}_{\geq 0}^2$, see [13], and for any non-negative function h on $\mathbb{Z}_{\geq 0}^2$ which is harmonic with respect to S killed outside $\mathbb{Z}_{\geq 0}^2$, there exists a positive measure μ_h on $\partial_M^{\mathbb{Z}_{\geq 0}^2} S$ such that

$$h(\,\cdot\,) = \int_{\partial_M^{\mathbb{Z}_{>0}^2} S} K_M(\,\cdot\,,\omega) d\mu_h(\omega)$$

One sees from (27) that a convenient way to compute the asymptotics of $K_M(\cdot, y)$ as y goes to infinity is to first study the behavior of G(x, y) when y goes to infinity in the cone.

We set $\Sigma = \mathbb{R}^2_{\geq 0} \cap \mathbb{S}^1$, and for $u \in \Sigma$, denote by $\phi(u)$ the unique solution of

(28)
$$\mathbb{E}\left(\exp\langle\phi(u), S(1)\rangle\right) = 1 \text{ and } \mathbb{E}\left(S(1)\exp\langle\phi(u), S(1)\rangle\right) = r_u u := \mu^u$$

for some $r_u > 0$. Remark that ϕ is bijective from Σ to $\overline{\mathcal{G}}_0$ (whose definition may be found in (16)), with

$$\phi^{-1}(a,b) = \frac{\nabla K(a,b)}{\|\nabla K(a,b)\|}$$

for $(a,b) \in \overline{\mathcal{G}_0}$. For $u \in \Sigma$, we then write \mathbb{P}_u for the probability measure on $\{S(n)\}_{n \ge 1}$ given by the transition probabilities

$$\mathbb{P}_u(S(1) \in A) = \mathbb{E}\left(\exp\langle\phi(u), S(1)\rangle \mathbf{1}_{S(1) \in A}\right)$$

for $A \subset \mathbb{Z}^2$, and we denote by \mathbb{E}_u the corresponding expectation. We then write Σ^u for the covariance matrix of S(1) under \mathbb{P}_u (recall that we denote by μ^u the drift of S(1) under \mathbb{P}_u , see (28)).

4.2. Statement of the main result. The main result of this section is the following theorem, which gives an asymptotics of G(x, y) when y goes to infinity along any direction in the quarter plane. Denote by $e_1 = (1, 0)$ and $e_2 = (0, 1)$ the two standard basis vectors.

Theorem 11. For any $x \in \mathbb{Z}_{>0}^2$ and $\{y(n)\}_{n \ge 1}$ with $|y(n)| \xrightarrow{n \to \infty} \infty$,

•
$$if \frac{y(n)}{|y(n)|} = u_n \text{ with } \lim u_n := u_0 \in \Sigma \setminus \{e_1, e_2\}, \text{ then}$$

$$\lim_{n \to \infty} \sqrt{|y(n)|} e^{-\langle \phi(u_n(y(n))), x - y(n) \rangle} G(x, y(n)) = A(u_0) \mathbb{P}_{u_0}(\tau_x = \infty);$$
• $if \frac{y(n)}{|y(n)|} = u_n \text{ with } \lim u_n = e_i, i \in \{1, 2\}, \text{ then, with } \bar{\imath} = 3 - i,$

$$\lim_{n \to \infty} \frac{|y(n)|^{3/2} e^{-\langle \phi(u(y(n))), x - y(n) \rangle}}{V(y_i(n))} G(x, y(n)) = A(e_i) \left(x_{\bar{\imath}} - \mathbb{E}_{e_i}(x_{\bar{\imath}} + S_{\bar{\imath}}(\tau_x))\right),$$

with $u \mapsto A(u)$ a continuous, positive function on Σ and V introduced in Proposition 15.

The proof of this theorem is postponed to Section 6. The main difficulty concerns the case of an asymptotic direction along the boundary axes, where the survival probability vanishes. The latter case is solved by first studying the Green function of the random walk in the half-planes $\mathbb{H}_1 := \mathbb{Z} \times \mathbb{Z}_{>0}$ and $\mathbb{H}_2 := \mathbb{Z}_{>0} \times \mathbb{Z}$. We achieve the latter in Section 5. To conclude this section, we collect a few useful estimates on the classical random walk on \mathbb{Z}^2 .

4.3. Preliminary estimates, and the Ney and Spitzer theorem. For all $u \in \Sigma$, we introduce the modified Green kernels

$$G_u(x,y) = \sum_{n \ge 1} \mathbb{P}_u(x + S(n) = y, \tau_x > n) \quad \text{and} \quad \widetilde{G}_u(x,y) = \sum_{n \ge 1} \mathbb{P}_u(x + S(n) = y)$$

For all $x, y \in \mathbb{Z}_{>0}^2$, we have

(29)
$$G_u(x,y) = e^{\langle \phi(u), y-x \rangle} G(x,y) \quad \text{and} \quad \widetilde{G}_u(x,y) = e^{\langle \phi(u), y-x \rangle} \widetilde{G}(x,y).$$

We recall the following result from Ney and Spitzer [25, Thm 2], using our notation $\mu^u = \mathbb{E}_u(S(1)).$

Theorem 12. There exists a continuous function $A : \Sigma \to \mathbb{R}_{>0}$ such that, as $t \to \infty$, uniformly on $u \in \Sigma$ and $x \in \mathbb{Z}^2$ with $|x| = o(t^{1/2})$,

$$\sqrt{t}\widetilde{G}_u(x,\lfloor t\mu^u \rfloor) \to A(u).$$

We should emphasize that the initial proof of Ney and Spitzer requires the random walk $\{S(n)\}_{n\geq 1}$ to be irreducible, and is valid only for fixed $x \in \mathbb{Z}^2$. However, the only reason for the first requirement is the use of the local large deviation limit theorem [25, Thm 2.1], which has since been proven for any random walk with finite generating function and whose support generates \mathbb{Z}^2 , see [27, Ch. 7, P10]. Likewise, the hypothesis of a fixed x can be relaxed to the condition $|x| = o(t^{1/2})$ (see the proof of Proposition 19 for a similar computation).

We will apply several times the following lemma, which gives a large deviation bound which is uniform on \mathbb{P}_u , $u \in \Sigma$.

Lemma 13. There exist constants $\eta, \eta' > 0$ such that for all $u \in \Sigma$, $n \ge 1$,

• for all $t \in (0, 1)$,

$$\mathbb{P}_u(|S(n) - n\mu^u| > tn) \leqslant 4\exp(-\eta t^2 n);$$

• for all $t \ge 1$,

$$\mathbb{P}_u(|S(n) - n\mu^u| > tn) \leq 4\exp(-\eta' tn).$$

Proof. Using Markov inequality, we get (denoting $S(n) = (S_1(n), S_2(n))$)

$$\mathbb{P}_u(|S_1(n) - n\mu_1^u| > tn) \leq 2\exp(-n\Lambda_{u,1}^*(t)),$$

where $\Lambda_{u,1}^*$ is the Legendre transform of the function

$$\alpha \in \mathbb{R} \mapsto \log \mathbb{E}_u(e^{\alpha(S_1(1)-\mu_1^u)}) = \log \mathbb{E}(e^{(\alpha+\phi(u)_1)S_1(1)}) - \alpha \mu_1^u.$$

An easy computation yields that

$$\Lambda_{u,1}^*(t) = \Lambda_1(t + \mu_1^u) - \phi(u)_1(t + \mu_1^u),$$

where Λ_1 is the Legendre transform of $\alpha \mapsto \log \mathbb{E}(e^{\alpha S_1(1)})$. Since $u \mapsto \mu_1^u$ and $u \mapsto \phi(u)_1$ are \mathcal{C}^2 , $u \mapsto \Lambda_{u,1}^*(t)$ is \mathcal{C}^2 . Since for all $u \in \Sigma$, we have $\Lambda_{u,1}^*(0) = (\Lambda_{u,1}^*)'(0) = 0$, and by convexity $(\Lambda_{u,1}^*)'(t) > 0$ for all $(u,t) \in \Sigma \times \mathbb{R}$, we deduce the existence of $\eta > 0$ such that for all $u \in \Sigma$ and all $t \leq 1$,

$$\Lambda_{u,1}^*(t) > \eta t^2.$$

Hence, uniformly on $u \in \Sigma$, for $t \leq 1$,

$$\mathbb{P}_u(|S_1(n) - n\mu_1^u| > tn) \leq 2\exp(-n\eta t^2).$$

Doing the same for $S_2(n)$ yields the result.

Likewise, by strict convexity there exists $\eta' > 0$ such that

$$\Lambda_{u,1}^*(t) > \eta' t$$

for $t \ge 1$, independently of u. The second result is then deduced as the first one.

Finally, we will use a uniform bound on the Green function \widetilde{G} on \mathbb{Z}^2 , namely, there exists C > 0 such that for all $x, y \in \mathbb{Z}^2$,

(30)
$$\widetilde{G}(x,y) \leqslant C.$$

The existence of such a bound is just a consequence of the transience of the walk $\{S(n)\}_{n \ge 1}$.

5. Asymptotics of the Green function on the half-space

The most delicate part of the description of the Martin boundary is related to the behavior of the Green function along the half-axes $\mathbb{R}_{>0}e_1$ and $\mathbb{R}_{>0}e_2$. This situation is dealt by first giving an asymptotics of the Green function on the half-planes \mathbb{H}_1 and \mathbb{H}_2 .

For $x, y \in \mathbb{H}_i$, denote by

(31)
$$\tau_x^i = \inf\{n \ge 0 : x + S(n) \notin \mathbb{H}_i\}$$

and introduce the Green kernels on the half-plane

$$\widehat{G}^i(x,y) = \sum_{n \geqslant 1} \mathbb{P}(x + S(n) = y, \tau^i_x > n) \quad \text{and} \quad \widehat{G}^i_u(x,y) = \sum_{n \geqslant 1} \mathbb{P}_u(x + S(n) = y, \tau^i_x > n).$$

The goal of this section is to prove the following result. For $z \in \mathbb{R}^2$, denote by z_1 (resp. z_2) its first (resp. second) coordinate.

Proposition 14. Let $\varepsilon, \eta > 0$ be small enough and $i \in \{1, 2\}$. As y goes to ∞ with $|y_i| = o(|y|^{1/2+\varepsilon})$, then uniformly on $x \in \mathbb{H}_i$ with $|x| = o(|y|^{1/2-\eta})$,

$$\widehat{G}^{i}(x,y) \sim \frac{B(u(y))x_{\overline{\imath}}}{|y|^{3/2}} V(y_{\overline{\imath}}) e^{\langle u(y), x-y \rangle},$$

where $u(y) = \frac{y}{|y|}, u \mapsto B(u)$ is a continuous, positive function on Σ and V will be introduced in Proposition 15.

We only prove it for i = 1, the other case being similar. The proof of Proposition 14 is decomposed in three cases, depending on the distance of the endpoint to the horizontal axis. To prove this result, we need two local limit theorems for the walk x + S(n)conditioned to stay in \mathbb{H}_1 .

The first local limit theorem concerns the case where the distance of the endpoint x + S(n) to the horizontal axis is similar to the fluctuation scale \sqrt{n} . The equivalence part of this result is given by [15, Lem. 3.2] when $x_2 = o(\sqrt{n})$ and by [15, Lem. 3.1 (c)] when $x_2 \ge c\sqrt{n}$ for some c > 0, whereas the uniform bound is given by [15, Lem. 2.3].

Proposition 15. There exists $\kappa > 0$ such that for all A > 0, as n goes to infinity, uniformly on $x, y \in \mathbb{H}_1$ with $|x| = o(\sqrt{n})$ and $|x - n\mu^{e_1}| \leq A\sqrt{n}$,

$$\mathbb{P}_{e_1}(x+S(n)=y,\tau_x>n) \sim \frac{\kappa x_2 V(y_2)}{2\pi \sqrt{\det(\Sigma^{e_1})}n^2} \exp\left(-\frac{\|C^{e_1}(y-n\mu^{e_1})\|^2}{2n}\right)$$

where V is a positive harmonic function with respect to $\{-S_2(n)\}_{n\geq 1}$ killed at the boundary. Moreover, there exists C > 0 such that for all $x, y \in \mathbb{H}_1$ with $|x| = o(\sqrt{n})$,

$$\mathbb{P}_{e_1}(x+S_n=y,\tau_x>n) \leqslant C \frac{x_2 V(y_2)}{n^2} \left(\exp\left(-\frac{\|C^u(y-n\mu^{e_1})\|^2}{2n}\right) + o\left(n^{-2}\right) \right).$$

Remark that in the latter proposition, C only depends on the behavior of $\frac{|x|}{\sqrt{n}}$. By [11, Lem. 13 (a)], we also have as z goes to ∞

$$(32) V(z) \sim z.$$

The second local limit theorem deals with the case where the distance of the endpoint x + S(n) to the horizontal axis is large compared to the fluctuation scale \sqrt{n} .

Proposition 16. For η small enough, as n goes to infinity, uniformly on $x, y \in \mathbb{H}_1$ with $|x| = o(\sqrt{n})$ and $u \in \Sigma$ with $n^{-1/2} = o(u_2)$ and $u_2 \leq n^{-1/2+\eta}$,

$$\mathbb{P}_{u}(x+S(n)=y,\tau_{x}>n) = \frac{c(u)x_{2}\mu_{2}^{u}}{\sum_{11}^{u}2\pi n\sqrt{\det(\Sigma^{u})}}\exp\left(-\frac{\|C^{u}(y-n\mu^{u})\|^{2}}{2n}\right) + o\left(\frac{x_{2}\mu_{2}^{u}}{n}\right),$$

with $o(\cdot)$ uniform on u and $x, y \in \mathbb{H}_1$. Moreover, there exists C > 0 such that for all $x, y \in \mathbb{H}_1$ with $|x| = o(\sqrt{n})$,

$$\mathbb{P}_{u}(x+S_{n}=y,\tau_{x}>n) \leqslant Cx_{2}\frac{\mu_{2}^{u}}{t} \left(\exp\left(-\frac{\|C^{u}(y-n\mu^{u})\|^{2}}{2n}\right) + o\left(n^{-2}\right)\right)$$

The proof of the latter proposition is very similar to the one of Proposition 15 and is given in Appendix A.

5.1. Asymptotics at mesoscopic distance. Since the support of the random walk S is $\{(i, j) \in \mathbb{Z}^2 : i, j \ge -1\}$ by our assumptions (ii), we have almost surely $S_2(1) \ge -1$. Such a random variable is called skip-free in the literature [7] and has interesting properties allowing to explicitly compute the survival probability.

For z > 0, denote by $\tau'_z = \inf\{n \ge 0 : z + S_2(n) = 0\}$, and remark that $\tau^1_x = \tau'_{x_2}$ for $x \in \mathbb{H}_1$, see (31). Then, the skip-free assumption yields that

- (SF1) if $\mu_2^u > 0$, $\mathbb{P}_u(\tau'_{x_2} < \infty) = \mathbb{P}_u(\tau'_1 < \infty)^{x_2} = c_u^{x_2}$, where $c_u \in (0, 1)$ is the unique real in (0, 1) such that $\mathbb{E}_u(c^{S_2(1)}) = 1$;
- (SF2) as μ_2^u goes to 0, one has $1 c_u \sim 2 \frac{\mu_2^u}{\Sigma_2^u}$.

We have the following first asymptotic result for the Green function on \mathbb{H}_1 , when the endpoint remains far from the horizontal axis.

Lemma 17. Let
$$\varepsilon > 0$$
. For any $x \in \mathbb{H}_1$, as $t \to \infty$, uniformly on $u \in \Sigma$ with $\mu_2^u \ge t^{-1/2+\varepsilon}$,
 $\sqrt{t}\widehat{G}_u(x, \lfloor t\mu^u \rfloor) \sim A(u)\mathbb{P}_u(\tau^1 = \infty) \sim B(u)x_2u_2.$

Proof. For $x \in \mathbb{H}_1$, we have $\tau_x^1 = \inf\{n \ge 0 : x_2 + S_2(n) \le 0\}$, see (31). Set $t_u = \lfloor t\mu^u \rfloor$. Then, for $n_t = \lfloor t^{1-\varepsilon} \rfloor$,

$$\begin{aligned} \widehat{G}_{u}(x,t_{u}) &= \sum_{n=1}^{n_{t}} \mathbb{P}_{u} \left(x + S(n) = t_{u}, \tau_{x}^{1} > n \right) + \mathbb{E}_{u} \left(G_{u}(x + S(n_{t}),t_{u}), \tau_{x_{2}}' > n_{t} \right) \\ &= \sum_{n=0}^{n_{t}} \mathbb{P}_{u}(x + S(n) = t_{u}, \tau_{x_{2}}' > n) + \mathbb{E}_{u} \left(\widetilde{G}_{u}(x + S(n_{t}),t_{u}), \tau_{x_{2}}' > n_{t} \right) \\ &- \mathbb{E}_{u} \left(\widetilde{G}_{u}(x + S(\tau_{x}),t_{u}), n_{t} < \tau_{x_{2}}' < \infty \right) \\ &=: S_{1} + S_{2} + S_{3}. \end{aligned}$$

By Lemma 13,

$$S_{1} \leq \sum_{n=1}^{n_{t}} \mathbb{P}_{u} \left(|S(n) - n\mu^{u}| \geq |\lfloor t\mu^{u} \rfloor - x - n\mu^{u}| \right) \leq \sum_{n=1}^{n_{t}} \mathbb{P}_{u} \left(|S(n) - n\mu^{u}| > \frac{|t\mu^{u}|}{2n} n \right)$$
$$\leq 4 \sum_{n=1}^{n_{t}} \exp\left(-\frac{\eta |t\mu^{u}|^{2}}{(2n)^{2}}\right) \leq n_{t} \exp\left(-\frac{\eta |t\mu^{u}|^{2}}{(2n_{t})^{2}}\right).$$

Hence, since $n_t \leq t^{1-\varepsilon}$,

$$S_1 = O\left(\exp(-ct^{2\varepsilon})\right)$$

for some c > 0. Then, set $\alpha = t^{1/2 - \varepsilon/4}$ and write

$$S_{2} = \mathbb{E}_{u} \Big(\widetilde{G}_{u}(x + S(n_{t}), t_{u}), |S(n_{t}) - n_{t}\mu^{u}| \leq \alpha, \tau_{x}^{1} > n_{t} \Big) \\ + \mathbb{E}_{u} \Big(\widetilde{G}_{u}(x + S(n_{t}), t_{u}), |S(n_{t}) - n_{t}\mu^{u}| > \alpha, \tau_{x}^{1} > n_{t} \Big) =: S_{21} + S_{22}.$$

First, by (30) and Lemma 13,

(33)
$$S_{22} \leqslant C\mathbb{P}_u(|S(n_t) - n_t\mu^u| > \alpha) \leqslant 4C \exp(-\eta(\alpha/n_t)^2 n_t) = O(\exp(-\eta t^{\varepsilon/2})).$$

Then, on the event $\{|S(n_t) - n_t \mu^u| \leq \alpha\}$, as t goes to infinity $\frac{|x + S(n_t) - n_t \mu|}{t} = o(t^{1/2-\varepsilon})$. Hence, by Theorem 12, uniformly on $x + S(n_t)$ on the event $\{|S(n_t)| \leq \alpha\}$,

$$\sqrt{t}\widetilde{G}_u(x+S(n_t),t_u) \xrightarrow{t\to\infty} A(u).$$

Hence, $S_{21} \sim_{t \to \infty} \frac{A(u)}{\sqrt{t}} \mathbb{P}_u(|S(n_t) - n_t \mu^u| \leq \alpha, \tau_x^1 > n_t)$ as t goes to ∞ . Split the latter probability as

$$\mathbb{P}_u(|S(n_t) - n_t \mu^u| \leq \alpha, \tau_x^1 > n_t) = \mathbb{P}_u(|S(n_t) - n_t \mu^u| \leq \alpha, n_t < \tau_x^1 < \infty) + \mathbb{P}_u(\tau_x^1 = \infty) - \mathbb{P}_u(|S(n_t) - n_t \mu^u| > \alpha, \tau_x^1 = \infty).$$

First,

$$\mathbb{P}_u\left(|S(n_t) - n_t\mu^u| > \alpha, \tau_x^1 = \infty\right) \leq \mathbb{P}_u\left(|S(n_t) - n_t\mu^u| > \alpha\right) = O(\exp(-\eta t^{\varepsilon/2}))$$

as in (33). Then, on the event $\{|S(n_t) - n_t \mu^u| \leq \alpha, \tau_x > n_t\}, v := x + S(n_t)$ satisfies

$$v_2 \ge x_2 + n_t \mu_2^u - \alpha \ge ct^{1-\varepsilon} \mu_2^u - t^{1/2-\varepsilon/4} \ge ct^{1/2}$$

for some constant c > 0, thanks to the condition $\mu_2^u \ge t^{-1/2+\varepsilon}$. Hence, by (SF1) and (SF2), on the event $\{|S(n_t) - n_t \mu^u| \le \alpha, \tau_x^1 > n_t\}$, we have

(34)
$$\mathbb{P}(\tau_{x+S(n_t)}^1 < \infty | |S(n_t) - n_t \mu^u| \le \alpha, \tau_x^1 > n_t) \le \left(1 - c_1 \frac{\mu_2^u}{\sum_{22}^u}\right)^{c_2 t^{1/2}} \le \exp(-ct^{\varepsilon})$$

for some constant c > 0, where we used again that $\mu_2^u \ge t^{-1/2+\varepsilon}$ in the last inequality. This implies that $\mathbb{P}_u(|S(n_t) - n_t \mu^u| \le \alpha, n_t < \tau_x^1 < \infty) = O(\exp(-ct^{\varepsilon}))$. Finally, by (SF1) and (SF2) with $\mu_2^u = o(1), \ \mu_2^u \ge t^{-1/2+\varepsilon}$,

$$\mathbb{P}_{u}(\tau'_{x_{2}} = \infty) = 1 - c_{u}^{x_{2}} \ge 1 - \left(1 - ct^{-1/2 + \varepsilon}\right)^{x_{2}} \ge c' x_{2} t^{-1/2 + \varepsilon}$$

for some constants c, c' depending on x, and we finally have $S_{21} \sim_{t \to \infty} \frac{A(u)}{\sqrt{t}} \mathbb{P}_u(\tau_x^1 = \infty)$ and then

$$S_2 \sim_{t \to \infty} \frac{A(u)}{\sqrt{t}} \mathbb{P}_u(\tau_x^1 = \infty).$$

Note that the latter term will be the main contribution to the asymptotics of $\widehat{G}_u(x, t_u)$. Finally, by (30),

$$S_{3} = \mathbb{E}_{u} \left(\widetilde{G}_{u}(x + S(\tau_{x}), t_{u}), n_{t} < \tau_{x}^{1} < \infty \right)$$

$$\leq C \mathbb{P}_{u} \left(n_{t} < \tau_{x}^{1} < \infty \right)$$

$$\leq C \left(\mathbb{P}_{u} \left(|S(n_{t}) - n_{t} \mu^{u}| \leq \alpha, n_{t} < \tau_{x}^{1} < \infty \right) + \mathbb{P}_{u} \left(|S(n_{t}) - n_{t} \mu| > \alpha \right) \right)$$

By (34), the first term is bounded by $\exp(-ct^{\varepsilon})$ and by Lemma 13 the second term is bounded by $\exp(-\eta t^{\varepsilon/2})$. Hence,

$$S_3 = o\left(\exp(-\eta t^{\varepsilon/2})\right).$$

Putting together the results on S_1 , S_2 and S_3 yields then

$$\widehat{G}_u(x, t_u) \sim_{t \to \infty} \frac{A(u)}{\sqrt{t}} \mathbb{P}_u(\tau_x^1 = \infty)$$

uniformly on u such that $u_2 \ge t^{-1/2+\varepsilon}$. The second equivalence in Lemma 17 is a consequence of (SF1) and (SF2).

5.2. Asymptotics of the Green function at microscopic distance. The proof essentially follows the original proof of Ney and Spitzer [25].

Proposition 18. Let $(\theta_t)_{t\geq 0}$ a positive function going to zero at ∞ and $\varepsilon > 0$. For $\eta > 0$ small enough, as t goes to ∞ , uniformly on $u \in \Sigma$ with $\frac{t^{-1/2}}{\theta_t} \leq u_2 \leq t^{-1/2+\eta}$ and $x \in \mathbb{H}_1$ with $|x| = o(t^{1/2-\varepsilon})$,

$$\sqrt{t}\widehat{G}_u(x,\lfloor t\mu^u \rfloor) \sim B(u)x_2u_2.$$

Proof. Write $y = \lfloor t\mu^u \rfloor$ and split $\widehat{G}_u(x, \lfloor t\mu^u \rfloor)$ as

 $\frac{t^{-1/2}}{\theta_t} \leqslant u_2 \leqslant t^{-1/2+\varepsilon},$

$$\begin{split} \widehat{G}_u(x, \lfloor t\mu^u \rfloor) &= \sum_{n=1}^{\lfloor t-A\sqrt{t} \rfloor - 1} \mathbb{P}_u\left(x + S(n) = y, \tau_x^1 > n\right) + \sum_{n=\lfloor t-A\sqrt{t} \rfloor}^{\lfloor t+A\sqrt{t} \rfloor} \mathbb{P}_u\left(x + S(n) = y, \tau_x^1 > n\right) \\ &+ \sum_{n=\lfloor t+A\sqrt{t} \rfloor + 1}^{\infty} \mathbb{P}_u\left(x + S(n) = y, \tau_x^1 > n\right) \\ &:= S_1 + S_2 + S_3. \end{split}$$

First, by the assumption on u_2 , $n^{-1/2} = o(u_2)$ and $u_2 \leq C n^{-1/2+\eta}$ for $\lfloor t - A\sqrt{t} \rfloor \leq n \leq \lfloor t + A\sqrt{t} \rfloor$, with $o(\cdot)$ and C only depending on A and θ_t . Hence, by Proposition 16, uniformly for $x, y \in \mathbb{H}_1$ with $|x| = o(\sqrt{n})$ and $\lfloor t - A\sqrt{t} \rfloor \leq n \leq \lfloor t + A\sqrt{t} \rfloor$ and $u \in \Sigma$ with

$$\mathbb{P}_u(x+S(n)=y,\tau_x>n) = \frac{c(u)x_2\mu_2^u}{\sum_{11}^u 2\pi n\sqrt{\det(\Sigma^u)}} \exp\left(-\frac{\|C^u(y-n\mu^u)\|^2}{2n}\right) + o\left(\frac{x_2\mu_2^u}{n}\right).$$

Applying the latter to S_2 yields

$$\sqrt{t}S_{2} = \sqrt{t} \left(\sum_{n=\lfloor t-A\sqrt{t} \rfloor}^{\lfloor t+A\sqrt{t} \rfloor} \frac{c(u)x_{2}\mu_{2}^{u}}{\Sigma_{11}^{u}2\pi n\sqrt{\det(\Sigma^{u})}} \exp\left(-\frac{\|C^{u}(y-n\mu^{u})\|^{2}}{2n}\right) + o\left(\frac{x_{2}\mu_{2}^{u}}{n}\right) \right) \\
= \sqrt{t} \frac{c_{u}x_{2}\mu_{2}^{u}}{\Sigma_{11}^{u}} \left(\sum_{n=\lfloor t-A\sqrt{t} \rfloor}^{\lfloor t+A\sqrt{t} \rfloor} \frac{1}{2\pi n\sqrt{\det(\Sigma^{u})}} \exp\left(-\frac{\|C^{u}(y-n\mu^{u})\|^{2}}{2n}\right) \right) + x_{2}\mu_{2}^{u}o(1).$$

By [25, Eq. (2.20)],

$$\sqrt{t} \sum_{n=\lfloor t-A\sqrt{t}\rfloor}^{\lfloor t+A\sqrt{t}\rfloor} \frac{1}{2\pi n\sqrt{\det(\Sigma^u)}} \exp\left(-\frac{\|C^u(y-n\mu^u)\|^2}{2n}\right) \xrightarrow{t\to\infty} \frac{1}{\sqrt{2\pi\det(\Sigma^u)}\|C^u\mu^u\|} + \varepsilon(A),$$

uniformly in $u \in \Sigma$, with $\varepsilon(A) \xrightarrow{A \to \infty} 0$ and $|x| = t^{1/2-\varepsilon}$. Hence, we also have

(35)
$$\sqrt{t}S_2 \sim_{t \to \infty} \frac{c_u x_2 \mu_2^u}{\sum_{11}^u \sqrt{2\pi \det(\Sigma^u)} \|C^u \mu^u\|} (1 + \varepsilon(A)).$$

Then, for $0 < \delta < 1$, split S_1 as

$$S_{1} = \sum_{n=1}^{\lfloor \delta t \rfloor} \mathbb{P}_{u} \left(x + S(n) = y, \tau_{x}^{1} > n \right) + \sum_{n=\lfloor \delta t \rfloor + 1}^{\lfloor t - A\sqrt{t} \rfloor - 1} \mathbb{P}_{u} \left(x + S(n) = y, \tau_{x}^{1} > n \right) =: S_{11} + S_{12}.$$

By Lemma 13, the first term is bounded above as

$$\frac{\sqrt{t}}{x_2\mu_2^u}S_{11} \leqslant \sqrt{t}\sum_{n=1}^{\lfloor \delta t \rfloor} \mathbb{P}_u\left(x+S(n)=y\right) \leqslant Ct^{3/2}\exp\left(-c(1-\delta)t\right) \xrightarrow{t\to\infty} 0$$

uniformly in $|x| < t^{1/2-\varepsilon}$, A > 0 and $u \in \Sigma$ (see [25, Eq. (2.22)]). By Proposition 16 with r = 3, the second term is bounded as

$$\frac{\sqrt{t}}{x_2\mu_2^u}S_{12} \leqslant \sqrt{t}\sum_{\substack{n=\lfloor\delta t\rfloor+1\\n=\lfloor\delta t\rfloor+1}}^{\lfloor t-A\sqrt{t}\rfloor-1} \left(\frac{Cx_2\mu_2}{n}\exp\left(-c^u\|y-n\mu^u\|^2\right)+o\left(n^{-2}\right)\right)$$
$$\leqslant \sqrt{t}\sum_{\substack{n=\lfloor\delta t\rfloor+1\\n=\lfloor\delta t\rfloor+1}}^{\lfloor t-A\sqrt{t}\rfloor-1} \left(\frac{Cx_2\mu_2}{n}\exp\left(-c^u\|y-n\mu^u\|^2\right)\right)+o\left(\frac{1}{t}\right).$$

By [25, Eq. (2.26)],

$$\limsup_{t \to \infty} \sqrt{t} \sum_{n = \lfloor \delta t \rfloor + 1}^{\lfloor t - A\sqrt{t} \rfloor - 1} \frac{1}{n} \exp\left(-c^u \|y - n\mu^u\|^2\right) = \varepsilon'(A)$$

uniformly on $u \in \Sigma$, with $\varepsilon'(A)$ going to 0 when A going to ∞ . Hence,

$$\limsup_{t\to\infty}\frac{\sqrt{t}}{x_2\mu_2^u}\sum_{n=\lfloor\delta t\rfloor+1}^{\lfloor t-A\sqrt{t}\rfloor-1}\mathbb{P}_u(x+S(n)=y,\tau_x^1>n)=\varepsilon'(A).$$

Finally,

(36)
$$\limsup_{t \to \infty} \frac{\sqrt{t}}{x_2 \mu_2^u} S_1 = \varepsilon'(A),$$

uniformly on $u \in \Sigma$ and $x \in \mathbb{H}_1$ with $|x| < t^{1/2-\varepsilon}$. One proves similarly that

(37)
$$\limsup_{t \to \infty} \frac{\sqrt{t}}{x_2 \mu_2^u} S_3 = \varepsilon'(A),$$

uniformly on $u \in \Sigma$ and $x \in \mathbb{H}_1$ with $|x| < t^{1/2-\varepsilon}$. Putting (35), (36) and (37) together, we get

$$1 - \varepsilon(A) - 2\varepsilon'(A) < \liminf \frac{\sqrt{t}\widehat{G}_u(x, \lfloor t\mu^u \rfloor)\Sigma_{11}^u \sqrt{2\pi \det(\Sigma^u)} \|C^u \mu^u\|}{c_u x_2 \mu_2^u} \\ \leqslant \limsup \frac{\sqrt{t}\widehat{G}_u(x, \lfloor t\mu^u \rfloor)\Sigma_{11}^u \sqrt{2\pi \det(\Sigma^u)} \|C^u \mu^u\|}{c_u x_2 \mu_2^u} < 1 + \varepsilon(A) + 2\varepsilon'(A)$$

for all A > 0. Letting A go to ∞ , we thus get

$$\lim_{t \to \infty} \frac{\sqrt{t} \widehat{G}_u(x, \lfloor t\mu^u \rfloor)}{x_2 \mu_2^u} = \frac{c_u}{\sum_{11}^u \sqrt{2\pi \det(\Sigma^u)} \|C^u \mu^u\|}$$

uniformly on $u \in \Sigma$ and $x \in \mathbb{H}_1$ with $|x| = o(t^{1/2-\varepsilon})$.

We give a similar asymptotic result along the horizontal axis.

Proposition 19. Let B > 0. Uniformly on $x \in \mathbb{H}_1$ with $|x| = o(t^{1/2-\varepsilon})$, $y = t\mu^{e_1} + t'e_2$ with $0 \leq t' \leq B\sqrt{t}$,

$$\widehat{G}_{e_1}(x,y) \sim \frac{1}{|y|^{3/2}} B(e_1) V(y_2') x_2 e^{-\langle \phi(u(y)) - \phi(e_1), y \rangle},$$

with V introduced in (32) and $B(e_1) > 0$.

Proof. The proof follows the lines of the one of Proposition 18, using Proposition 15 instead of Proposition 16. The only different step is the computation of

$$\sqrt{t} \left(\sum_{n=\lfloor t-A\sqrt{t}\rfloor}^{\lfloor t+A\sqrt{t}\rfloor} \frac{1}{2\pi n\sqrt{\det(\Sigma^u)}} \exp\left(-\frac{\|C^u(\lfloor t\mu^u \rfloor - n\mu^u)\|^2}{2n}\right) \right)$$

which has to be replaced by

$$\sqrt{t} \left(\sum_{n=\lfloor t-A\sqrt{t}\rfloor}^{\lfloor t+A\sqrt{t}\rfloor} \frac{1}{2\pi n^2 \sqrt{\det(\Sigma^{e_1})}} \exp\left(-\frac{\|C^{e_1}(\lfloor t\mu^{e_1}+t'e_2\rfloor-n\mu^{e_1})\|^2}{2n}\right) \right) := S_A$$

As t goes to infinity and for $\lfloor t - A\sqrt{t} \rfloor \leq n \leq \lfloor t + A\sqrt{t} \rfloor$, $n \sim t$, and thus

$$S_A \sim \frac{1}{\sqrt{t}} \left(\sum_{n=\lfloor t-A\sqrt{t} \rfloor}^{\lfloor t+A\sqrt{t} \rfloor} \frac{1}{2\pi n\sqrt{\det(\Sigma^{e_1})}} \exp\left(-\frac{\|C^{e_1}(\lfloor t\mu^{e_1}+t'e_2 \rfloor - n\mu^{e_1})\|^2}{2n}\right) \right).$$

By [25], uniformly on $t' \leq B\sqrt{t}$,

$$\lim_{t \to \infty} \sum_{n = \lfloor t - A\sqrt{t} \rfloor}^{\lfloor t + A\sqrt{t} \rfloor} \frac{1}{2\pi n\sqrt{\det(\Sigma^{e_1})}} \exp\left(-\frac{\|C^{e_1}(\lfloor t\mu^{e_1} + t'e_2 \rfloor - n\mu^{e_1})\|^2}{2n}\right) \\ \sim A(u(y)) \exp(-\langle \phi(u(y)) - \phi(e_1), y \rangle) + \varepsilon(A),$$

with $\varepsilon(A)$ going to 0 as A goes to infinity. Hence,

$$\lim_{A \to \infty} \lim_{t \to \infty} \sqrt{t} S_A = A(u(y)) \exp(-\langle \phi(u(y)) - \phi(e_1), y \rangle)$$

Concluding the proof as the one of Proposition 18, we get

$$\widehat{G}_{e_2}(x,y) \sim_{t \to \infty} \frac{B(e_2)}{|y|^{3/2}} V(y_2) x_2 e^{-\langle \phi(u(y)) - \phi(e_1), y \rangle},$$

with $B(e_2) = \kappa \widetilde{A}(e_2)$.

Proof of Proposition 14. When $\sqrt{|y|} = o(y_2)$, the statement of Proposition 14 is obtained by putting together Lemma 17 and Proposition 18 with u = u(y) and $t = \frac{|y|}{|\mu^{u(y)}|}$, using that $\mu_2^{u(y)} \sim \frac{y_2}{|y|}$ and $V(z) \sim z$ as z goes to infinity. When $y_2 \leq B|y_1|^{1/2}$ for some B > 0, use Proposition 19 with $t = \frac{y_1}{|\mu^{e_1}|} \sim \frac{|y|}{|\mu^{e_1}|}$ to get

$$\widehat{G}(x,y) \sim \frac{B(e_1)x_2V(y_2)}{|y|^{3/2}} e^{\langle \phi(e_1), y \rangle - \langle \phi(u(y)), y \rangle}$$

with $B(e_1) > 0$. The result is deduced provided we prove that

$$\langle \phi(e_1), y \rangle = \langle \phi(u(y)), x \rangle + o(1)$$

uniformly on $y = o(|x|^{1/2-\varepsilon})$. As y goes to infinity, $u(y) = e_1 + \frac{y_2}{|y|}e_2 + o(\frac{y_2}{|y|})$. Since ϕ is \mathcal{C}^2 , $K(\phi(u)) = 1$ and $\nabla K \circ \phi(u) \in \mathbb{R}u$. For $\delta > 0$ small enough with $u + \delta \in \Sigma$, we have $\phi(u + \delta) = \phi(u) + u^{\perp}h_u(\delta) + O(|\delta|^2)$ for some linear functional h_u . Hence,

$$\phi(u(y)) = \phi(e_1) + \alpha \frac{y_2}{|y|} e_2 + O\left(\frac{y_2}{|y|}\right)^2.$$

Therefore, for $x = o(|y|^{1/2-\varepsilon})$,

$$\langle \phi(u(y)), x \rangle = \langle \phi(e_1), y \rangle + o\left(\frac{y_2}{|y|^{1/2+\varepsilon}}\right) = \langle \phi(e_1), y \rangle + o(1).$$

The result is then deduced.

6. Identification of the Martin Boundary

We can now describe the Martin boundary $\partial_{\mathbb{Z}_{>0}^{M}}^{M} S$. We first prove Theorem 11, and then deduce from the latter the following description:

Corollary 20. The Martin boundary $\partial_M^{\mathbb{Z}_{\geq 0}^2} S$ of $\{S(n)\}_{n \geq 0}$ killed outside of the quarter plane $\mathbb{Z}_{\geq 0}^2$ is homeomorphic to $\overline{\mathcal{G}_0}$ through the map

$$(a_0, b_0) \in \overline{\mathcal{G}_0} \mapsto \frac{h_{(a_0, b_0)}}{h_{(a_0, b_0)}(1, 1)} \in \partial_M^{\mathbb{Z}_{>0}^2} S.$$

More precisely, for $(a_0, b_0) \in \mathcal{G}_0$, writing $u = \frac{\nabla K(a_0, b_0)}{\|\nabla K(a_0, b_0)\|}$,

$$h_{(a_0,b_0)}(x) = e^{\langle (a_0,b_0),x \rangle} \mathbb{P}_u(\tau_x = \infty) = e^{\langle (a_0,b_0),x \rangle} - \mathbb{E}\left(e^{\langle (a_0,b_0),x + S(\tau_x) \rangle}, \tau_x < \infty\right)$$

and if $(a_0, b_0) \in \partial \mathcal{G}_0$, with $e_i = \frac{\nabla K(a_0, b_0)}{\|\nabla K(a_0, b_0)\|}, i \in \{1, 2\},\$

$$h_{(a_0,b_0)}(x) = x_{\overline{\imath}} e^{\langle (a_0,b_0),x\rangle} - \mathbb{E}\left(e^{\langle (a_0,b_0),x+S(\tau_x)\rangle}(x_{\overline{\imath}} + S_{\overline{\imath}}(\tau_x))\right).$$

Proof of Theorem 11. We only prove the first statement and sketch the proof of the second one, which is similar to the first proof.

Let $V(u_0)$ be a neighborhood of $u_0 \in \Sigma$ such that $d(V(u_0), \partial \Sigma) > 0$. Set $t_n = \frac{|y(n)|}{|\mu^{u(y(n))}|}$ and $u_n = \frac{y(n)}{|y(n)|}$. Then, for $m_n \ge 1$ to adjust later,

$$G_{u_n}(x, y(n)) = \sum_{m=1}^{m_n} \mathbb{P}_{u_n}(x + S(m) = y(n), \tau_x > m) + \mathbb{E}_{u_n} (G_{u_n}(x + S(m_n), y(n)), \tau_x > m_n)$$

$$= \sum_{m=0}^{m_n} \mathbb{P}_{u_n}(x + S(m) = y(n), \tau_x > m) + \mathbb{E}_{u_n} \left(\widetilde{G}_u(x + S(m_n), y(n)), \tau_x > m_n \right)$$

(38)

$$= \mathbb{E}_{u_n} \left(\widetilde{G}_u(x + S(\tau_x), y(n)), m_n < \tau_x < \infty \right)$$

$$=: S_1 + S_2 + S_3.$$

By the second part of Lemma 13, as long as $m_n < t_n/3$ and for |y(n)| large enough

$$S_{1} \leqslant \sum_{n=1}^{m_{n}} \mathbb{P}_{u_{n}} \left(|S(m) - m\mu^{u(n)}| \geqslant \left| \lfloor t_{n}\mu^{u(n)} \rfloor - x - m\mu^{u(n)} \right| \right)$$
$$\leqslant \sum_{n=1}^{m_{n}} \mathbb{P}_{u_{n}} \left(|S(m) - m\mu^{u(n)}| > \frac{|t_{n}\mu^{u(n)}|}{2m} m \right)$$
$$\leqslant \sum_{n=1}^{m_{n}} \exp \left(-\frac{\eta'|t\mu^{u(n)}|}{2m} \right) \leqslant m_{n} \exp \left(-\frac{\eta'|t_{n}\mu^{u(n)}|}{2m_{n}} \right).$$

Hence, as $m_n = o(t_n^{1/4})$,

(39)
$$S_1 = o(\exp(-\sqrt{t_n})).$$

Then, write

$$S_{2} = \mathbb{E}_{u_{n}} \left(\widetilde{G}_{u_{n}}(x + S(m_{n}), y(n)), |S(m_{n}) - m_{n}\mu^{u(n)}| \leq 2m_{n}, \tau_{x} > m_{n} \right) \\ + \mathbb{E}_{u_{n}} \left(\widetilde{G}_{u_{n}}(x + S(m_{n}), y(n)), |S(m_{n}) - m_{n}\mu^{u(n)}| > 2m_{n}, \tau_{x} > m_{n} \right) \\ = S_{21} + S_{22}.$$

First, by (30) and Lemma 13,

$$S_{22} \leqslant C\mathbb{P}_{u_n}(|S(m_n) - m_n \mu^{u(n)}| > 2m_n) \leqslant C \exp(-2\eta' m_n).$$

Then, set $\widetilde{u} = \frac{y(n)-(x+S(m_n))}{|y(n)-(x+S(m_n))|} \in \mathbb{S}^1$. There exists t_0 such that on the event $\{|S(m_n) - n\mu^u| \leq 2m_n\}$, for $t_n \geq t_0$, $\widetilde{u} \in V(u_0)$. Then, by Theorem 12, as t_n goes to infinity with $m_n = o(t_n^{1/4})$, uniformly on $x + S(m_n)$ on the event $\{|S(m_n)| \leq 2m_n\}$ and using the fact that $u_n \to u_0$,

$$\sqrt{t}\widetilde{G}_{u_n}(x+S(m_n),y(n)) \xrightarrow{t\to\infty} A(u_0).$$

Hence,

$$S_{21} \sim_{n \to \infty} \mathbb{E}_{u_n} \left(\frac{A(u_0)}{\sqrt{t_n}} \mathbf{1}_{|S(m_n) - m_n \mu^{u(n)}| \leq 2m_n, \tau_x > m_n} \right)$$
$$= \frac{A(u_0)}{\sqrt{t_n}} \mathbb{P}_{u_n} \left(|S(m_n) - m_n \mu^{u(n)}| \leq 2m_n, \tau_x > m_n \right).$$

Since

$$\mathbb{P}_{u_n}\left(|S(m_n) - m_n \mu^{u(n)}| \le 2m_n, \tau_x > m_n\right) \\ = \mathbb{P}_{u_n}\left(\tau_x > m_n\right) - \mathbb{P}_{u_n}\left(|S(m_n) - m_n \mu^{u(n)}| > 2m_n, \tau_x > m_n\right),$$

with

$$\mathbb{P}_{u_n}(\tau_x > m_n) \sim_{n \to \infty} \mathbb{P}_{u_n}(\tau_x = \infty) \sim_{n \to \infty} \mathbb{P}_{u_0}(\tau_x = \infty)$$

and $\mathbb{P}_{u_n}(|S(m_n) - m_n \mu^{u(n)}| > 2m_n, \tau_x > m_n) \leq \exp(-2\eta' m_n)$ by Lemma 13,

$$S_{21} \sim_{n \to \infty} \frac{A(u) \mathbb{P}_{u_0}(\tau_x = \infty)}{\sqrt{t_n}},$$

and as long as $m_n = o(t_n^{1/4})$, and $\log(t_n) = o(m_n)$,

(40)
$$S_2 \sim_{t \to \infty} \frac{A(u_0) \mathbb{P}_{u_0}(\tau_x = \infty)}{\sqrt{t}}.$$

As in the proof of Lemma 17, the latter term will be the main contribution to the asymptotics of $G_{u_n}(x, y(n))$. Finally, by (30),

$$S_{3} = \mathbb{E}_{u_{n}} \left[\widetilde{G}_{u_{n}}(x + S_{\tau_{x}}, y(n)), m_{n} < \tau_{x} < \infty \right]$$

$$\leq C \mathbb{P}_{u_{n}} \left[m_{n} < \tau_{x} < \infty \right]$$

$$\leq C \left(\mathbb{P}_{u_{n}} \left(|S(m_{n}) - m_{n}\mu^{u(n)}| \leq cm_{n}, m_{n} < \tau_{x} < \infty \right) + \mathbb{P}_{u_{n}} \left(|S(m_{n}) - m_{n}\mu| > cm_{n} \right) \right),$$

for c small enough so that $\inf_{u \in V(u_0)} d(\mu^{u(n)}, \partial \mathbb{R}^2_{>0}) := d > c$. On the one hand, it follows from Lemma 13 that there exists η' such that

$$\mathbb{P}_{u_n}(|S(m_n) - m_n \mu^{u(n)}| > cm_n) \leqslant \exp(-\eta' m_n)$$

for all $m_n \ge 1$ and $u \in V(u_0)$. On the other hand,

$$\mathbb{P}_{u_n}(|S(m_n) - m_n \mu^{u(n)}| \le cm_n, m_n < \tau_x < \infty) \le \sup_{\substack{y \in B(x + S(m_n), cm_n)}} \mathbb{P}_{u_n}(\tau_y < \infty)$$
$$\le \sup_{\substack{y \in B(x + S(m_n), cm_n)}} \mathbb{P}_{u_n}(\tau_y^i < \infty),$$

for $i \in \{1, 2\}$, where $\tau_y^i = \inf\{n \ge 0 : x_i + S_i(n) \le 0\}$. Since S is skip-free and there exists $\mu_i > 0$ such $\mu_i^u \ge \mu_i$ for all $u \in V(u_0)$, by (SF1) and (SF2) there exists $c_{u_0} > 0$ independent of $u \in V(u_0)$ such that $\mathbb{P}_{u_n}(\tau_y^i < \infty) \le c_{u_0}^y$. Hence, by the choice of c we have

(41)
$$\sup_{y \in B(x+S(m_n),c_{m_n})} \mathbb{P}_{u_n}(\tau_y < \infty) \leqslant \exp(-c_{u_0}(d-c)m_n).$$

Finally, there exists c > 0 independent of $u \in V(u_0)$ such that $S_3 \leq \exp(-cm_n)$, and as long as $\log(t_n) = o(m_n)$,

$$(42) S_3 = o(\sqrt{t_n}).$$

Putting (39), (40) and (42) together yields

$$\sqrt{t_n}G_{u_n}(x,y(n)) \xrightarrow[t \to \infty]{} A(u_0)\mathbb{P}_{u_0}(\tau_x = \infty),$$

and setting $B(u_0) = A(u_0)|\mu^{u_0}|$ yields the desired limit for G_{u_n} , since $t_n = \frac{|y(n)|}{|\mu^{u_n}|} \sim_{n \to \infty} \frac{|y(n)|}{|\mu^{u_0}|}$. Using (29) yields then the final result.

The proof of the second statement is similar. The unique change is to use Proposition 14 instead of Theorem 12, and the renewal relation

$$G(x,y) = \widehat{G}_i(x,y) - \mathbb{E}(\widehat{G}(x+S(\tau_x^{\overline{i}}),y),\tau_x^{\overline{i}} < \tau_x^i)$$

in (38). The main term S_{21} is then asymptotically equivalent to

$$\frac{A(e_i)V(y_2(n))f(y_2'/\sqrt{|y(n)|})}{|y(n)|^{3/2}}\mathbb{E}_{e_i}(x_i+S_i(m_n),\tau_x>m_n).$$

Then, using that $x_{\bar{i}}$ is harmonic for S killed at τ^i under \mathbb{P}_{e_i} (because S is skip-free),

$$\begin{aligned} &\mathbb{E}_{e_i}(x_{\overline{\imath}} + S_{\overline{\imath}}(m_n), \tau_x > m_n) \\ &= \mathbb{E}_{e_i}(x_{\overline{\imath}} + S_{\overline{\imath}}(m_n), \tau_x^i > m_n) - \mathbb{E}_{e_i}(x_{\overline{\imath}} + S_{\overline{\imath}}(m_n), \tau_x^i > m_n \ge \tau^{\overline{\imath}}) \\ &= x_{\overline{\imath}} - \mathbb{E}_{e_i}\left(\mathbb{E}_{e_i}\left(x_{\overline{\imath}} + S_{\overline{\imath}}(\tau^{\overline{\imath}}) + \widetilde{S}_{\overline{\imath}}(m_n - \tau^{\overline{\imath}}), \tau_{x+S(\tau^{\overline{\imath}}_x)}^i > m_n - \tau^{\overline{\imath}}_x\right), \tau_x^{\overline{\imath}} \le m_n, \tau^{\overline{\imath}} < \tau_x^i)\right), \end{aligned}$$

where \widetilde{S} has the same law of S and is independent of $\tau^{\overline{i}}$. By harmonicity of $x_{\overline{i}}$ for S killed at τ^i , we get

$$\mathbb{E}_{e_i}\left(x_{\bar{\imath}} + S_{\bar{\imath}}(\tau^{\bar{\imath}})_x + \widetilde{S}_{\bar{\imath}}(m_n - \tau_x^{\bar{\imath}}), \tau_{x_{\bar{\imath}} + S_{\bar{\imath}}(\tau_x^{\bar{\imath}})}^i > m_n - \tau_x^{\bar{\imath}}\right) = x_{\bar{\imath}} + S_{\bar{\imath}}(\tau_x^{\bar{\imath}}),$$

so that

$$\mathbb{E}_{e_i}(x_{\bar{\imath}} + S_{\bar{\imath}}(m_n), \tau_x > m_n) = x_{\bar{\imath}} - \mathbb{E}_{e_i}\left(x_{\bar{\imath}} + S_{\bar{\imath}}(\tau^{\bar{\imath}}), \tau^{\bar{\imath}} \leqslant m_n, \tau_x^{\bar{\imath}} < \tau_x^i\right).$$

Since $\mathbb{P}(m_n < \tau_x^{\overline{i}} < \infty) = o(\exp(-t_n^{\varepsilon}))$ for some $\varepsilon > 0$ as in (41), we have

$$\mathbb{E}_{e_i}(x_{\overline{\imath}} + S_{\overline{\imath}}(m_n), \tau_x > m_n) \sim x_{\overline{\imath}} - \mathbb{E}_{e_i}\left(x_{\overline{\imath}} + S_{\overline{\imath}}(\tau_x^{\overline{\imath}}), \tau_x^{\overline{\imath}} < \tau_x^i\right) = x_{\overline{\imath}} - \mathbb{E}_{e_i}\left(x_{\overline{\imath}} + S_{\overline{\imath}}(\tau_x)\right) \\ = -\mathbb{E}_{e_i}\left(S_{\overline{\imath}}(\tau_x)\right),$$

where we used that $x_{\bar{\imath}} + S_{\bar{\imath}}(\tau_x^i) = 0$ because S is skip-free.

Proof of Corollary 20. By Theorem 11, if $\{y(n)\}_{n\geq 0}$ is such that $|y(n)| \to \infty$ with $\frac{y(n)}{|y(n)|} \to u \in \Sigma$, then if $u \notin \{e_1, e_2\}$, for $x \in \mathbb{Z}^2_{>0}$ we have

$$K(x, y(n)) \to e^{\langle \phi(u), x - x_0 \rangle} \frac{\mathbb{P}_{u_n}(\tau_x = \infty)}{\mathbb{P}_{u_n}(\tau_{x_0} = \infty)} = C(u)e^{\langle \phi(u), x \rangle} \mathbb{P}_{u_n}(\tau_x = \infty)$$

$$(43) = C(u)\left(e^{\langle \phi(u), x \rangle} - \mathbb{E}\left(e^{\langle \phi(u), x + S(\tau) \rangle}, \tau < \infty\right)\right)$$

$$:= \bar{h}_{\phi(u)}(x),$$

for some C(u) > 0 and if $u = e_i, i \in \{1, 2\},\$

$$(44) K(x,y(n)) \to e^{\langle \phi(e_i), x-x_0 \rangle} \frac{x_i - \mathbb{E}_{e_i}(x_i + S_i(\tau_x))}{x_{0i} - \mathbb{E}_{e_i}(x_{0i} + S_i(\tau_{x_0}))} = C(e_i) \left(x_i e^{\langle \phi(e_i), x \rangle} - \mathbb{E} \left(e^{\langle \phi(e_i), x + S(\tau) \rangle}(x_i + S_i(\tau_x)) \right) \right) := \bar{h}_{\phi(e_i)}(x),$$

for some $C(e_i) > 0$. Remark that as $d(x, \partial \mathbb{Z}_{>0}) \to \infty$, $\mathbb{P}_{u_n}(\tau_x = \infty) \to 1$ and $\mathbb{E}_{e_i}(x_{\bar{\imath}} + S_{\bar{\imath}}(\tau_x^i), \tau_x^{\bar{\imath}} > \tau_x^i) \to \mathbb{E}_{e_i}(x_{\bar{\imath}} + S_{\bar{\imath}}(\tau_x^i)) = x_i$, so that if $u \notin \{e_1, e_2\}$,

$$\bar{h}_{\phi(u)}(x) \sim_{d(x,\partial\mathbb{Z}_{>0})\to\infty} e^{\langle\phi(u),x\rangle}$$

and if $u = e_i, i \in \{1, 2\},\$

$$\bar{h}_{\phi(e_i)}(x) \sim_{d(x,\partial\mathbb{Z}_{>0})\to\infty} x_{\bar{\imath}} e^{\langle \phi(e_i), x \rangle}$$

Since $u \mapsto \phi(u)$ is injective, we deduce that $u \mapsto \bar{h}_{\phi(u)}$ is injective. Hence, $\Phi : u \mapsto \bar{h}_{\phi(u)}$ is a bijection from Σ to $\partial_M^{\mathbb{Z}_{\geq 0}^2}S$. Remark that Φ is continuous: this is clear from (43) on $\Sigma \setminus \{e_1, e_2\}$, and (44) for e_1 is obtained as a limit of (43) as u goes to e_1 (the derivative appearing because of the term C(u)). Since Σ is compact and Φ is a continuous bijection, Φ is a homeomorphism from Σ to $\partial_M^{\mathbb{Z}_{\geq 0}^2}S$.

It remains to prove that $\frac{h_{\phi(u)}}{h_{\phi(u)}(1,1)} = \bar{h}_{\phi(u)}$. Let us first suppose that $u \in \Sigma \setminus \{e_1, e_2\}$. The function $h_{\phi(u)}$ is harmonic in $\mathbb{Z}_{>0}^2$, so that $h_{\phi(u)} = \int_{\Sigma} \bar{h}_{\phi(w)} dm_u(w)$ for some positive measure m_u on Σ . We have $h_{\phi(u)}(x) \sim e^{\langle \phi(u), x \rangle}$ as $d(x, \partial \mathbb{Z}_{>0}) \to \infty$. Hence, since $\bar{h}_{\phi(w)}(x) \sim e^{\langle \phi(w), x \rangle}$ (resp. $\bar{h}_{\phi(e_i)}(x) \sim x_{\bar{i}} e^{\langle \phi(e_i), x \rangle}$ as $d(x, \partial \mathbb{Z}_{>0}) \to \infty$ for $w \in \Sigma \setminus \{e_1, e_2\}$ (resp. $w = e_i, i \in \{1, 2\}$) we must have $m_u = c\delta_u$ for come constant c. Since $\frac{h_{\phi(u)}(1,1)}{h_{\phi(u)}(1,1)} = \bar{h}_{\phi(u)}(1,1) = 1, c = 1$ and thus

$$\frac{h_{\phi(u)}}{h_{\phi(u)}(1,1)} = \bar{h}_{\phi(u)}$$

Then for $i \in \{1, 2\}$, by continuity of Φ and Proposition 6,

$$\frac{h_{\phi(e_i)}}{h_{\phi(e_i)}(1,1)} = \lim_{u \to e_i} \frac{h_{\phi(u)}}{h_{\phi(u)}(1,1)} = \lim_{u \to e_i} \bar{h}_{\phi(u)} = \bar{h}_{\phi(e_i)}.$$

Since ϕ is a homeomorphism from Σ to $\overline{\mathcal{G}_0}$ with $\phi^{-1}(a_0, b_0) = \frac{\nabla K(a_0, b_0)}{\|\nabla K(a_0, b_0)\|}$, we get the statement of the corollary.

APPENDIX A. PROOF OF PROPOSITION 16

The proof of Proposition 16 is very similar to the one of Theorem 5 in [11] which provides a local limit theorem for zero-drift random walks in cones (the cone being a half-space for us) : the two differences are the presence of a small drift in our case and the fact that we are dealing with walks with small negative jumps in the vertical direction, which greatly simplifies the proof. The first step is to get good estimates on the heat kernel of the Brownian motion with drift inside $\mathbb{R} \times \mathbb{R}_{>0}$.

Lemma 21. Let $(B_t)_{t\geq 0}$ be a Brownian motion with drift μ and covariance Σ . Denote by τ_x^B the exit time from $\mathbb{R} \times \mathbb{R}_{\geq 0}$ for $x + B_t$. Then, writing $C = \Sigma^{-1/2}$ and $K^B(x, y)dy = d\mathbb{P}(x + B_t = y, \tau_x^B > t)$, we have for $x, y \in \mathbb{R} \times \mathbb{R}_{\geq 0}$

$$K_t^B(x,y) = \left(1 - e^{-\frac{2x_2y_2}{t\Sigma_{11}}}\right) \frac{\exp\left(-\frac{\|C(y-x-t\mu)\|^2}{2t}\right)}{2\pi t \det(\Sigma)}$$

In particular, for all $x, y \in \mathbb{R} \times \mathbb{R}_{\geq 0}$

$$K_t^B(x,y) \leqslant \frac{2x_2y_2}{t\Sigma_{11}} \frac{\exp\left(-\frac{\|C(y-x-t\mu)\|^2}{2t}\right)}{2\pi t \det(\Sigma)},$$

and as $\frac{x_2y_2}{\Sigma_{11}} \leq \theta_t t$ with θ_t going to zero,

$$K_t^B(x,y) \sim \frac{2x_2y_2}{t\Sigma_{11}} \frac{\exp\left(-\frac{\|C(y-x-t\mu)\|^2}{2t}\right)}{2\pi t \det(\Sigma)},$$

with ~ only depending on θ_t .

Proof. This lemma is a consequence of Girsanov theorem and the reflection formula. For $v \in \mathbb{R}^2$, denote by s_v the orthogonal symmetry with respect to a one-dimensional vector space generated by v, so that $s_v(x) = x - 2 \frac{\langle x, v^{\perp} \rangle}{\langle v^{\perp}, v^{\perp} \rangle} v^{\perp}$ for any $v^{\perp} \neq 0$ with $\langle v^{\perp}, v \rangle = 0$.

Recall that if W is a standard two-dimensional Brownian motion with zero drift and $\tau_x^v = \inf\{t \ge 0 : x + W_t \in \mathbb{R}v\}$, the reflection formula yields

$$\begin{split} d\mathbb{P}(x+W_t &= y, \tau_x^v > t) \\ &= \frac{1}{2\pi t} \left(\exp\left(-\frac{\|y-x\|^2}{2t}\right) - \exp\left(-\frac{\|s_v(y)-x\|^2}{2t}\right) \right) \\ &= \frac{1}{2\pi t} \left(\exp\left(-\frac{\|y-x\|^2}{2t}\right) - \exp\left(-\frac{\|y-x-2\frac{\langle y,v^{\perp}\rangle}{\langle v^{\perp},v^{\perp}\rangle}v^{\perp}\|^2}{2t}\right) \right) \\ &= \frac{1}{2\pi t} \left(\exp\left(-\frac{\|y-x\|^2}{2t}\right) - \exp\left(-\frac{\|y-x\|^2 - 4\frac{\langle y-x,v^{\perp}\rangle \cdot \langle y,v^{\perp}\rangle}{\langle v^{\perp},v^{\perp}\rangle} + 4\frac{\langle y,v^{\perp}\rangle^2}{\langle v^{\perp},v^{\perp}\rangle} \right) \right) \\ &= \frac{1 - e^{-\frac{2\langle y,v^{\perp}\rangle \langle x,v^{\perp}\rangle}{t\langle v^{\perp},v^{\perp}\rangle}}}{2\pi t} \exp\left(-\frac{\|y-x\|^2}{2t}\right). \end{split}$$

Write $\widehat{B}_t = CB_t$ and $\widetilde{B}_t = \widehat{B}_t - tC\mu$, so that \widehat{B} is a Brownian motion with covariance identity and drift $C\mu$ and \widetilde{B} is a standard Brownian motion. Using the change of variable $u \mapsto Cu$ and Girsanov theorem applied to \widetilde{B} , we get

$$\begin{split} K^B_t(x,y) &= d\mathbb{P}(x+B_t=y,\tau^B_x>t) \\ &= (\det C)d\mathbb{P}(Cx+\widehat{B}_t=Cy,\tau^{Ce_2}_x>t) \\ &= (\det C)e^{\langle C\mu,C(x-y)\rangle - \frac{t\|C\mu\|^2}{2}}d\mathbb{P}(Cx+\widetilde{B}_t=Cy,\tau^{Ce_1}_x>t). \end{split}$$

Then, since $\langle \Sigma^{1/2} e_2, C e_1 \rangle = 0$, we can apply the previous computation to get

$$\begin{split} K_t^B(x,y) = &(\det C)e^{C\mu \cdot (x-y) - \frac{t\|C\mu\|^2}{2}} \frac{1 - e^{-\frac{2(Cy,C^{-1}e_2)\langle x,C^{-1}e_2\rangle}{te_1^T \Sigma e_1}}}{2\pi t} \exp\left(-\frac{\|Cy - Cx\|^2}{2t}\right) \\ = &\frac{1 - e^{-\frac{2y_2x_2}{t\Sigma_{11}}}}{2\pi t\sqrt{\det \Sigma}} \exp\left(-\frac{\|Cy - Cx\|^2 - 2t\langle C(y-x), C\mu\rangle + t^2\langle C\mu, C\mu\rangle}{2t}\right) \\ = &\frac{1 - e^{-\frac{2y_2x_2}{t\Sigma_{11}}}}{2\pi t\sqrt{\det \Sigma}} \exp\left(-\frac{\|C(y - x - t\mu)\|^2}{2t}\right). \end{split}$$

The rest of the lemma is deduced by using the formula $1 - e^{-u} \leq u$ for $u \in \mathbb{R}$ and $1 - e^{-u} \sim u$ as u goes to 0.

We next turn to the proof of Proposition 16.

Proof of Proposition 16. The proof of this proposition is exactly the same as the one of [11, Thm 5] and [15, Prop. 2.4]. We introduce the main steps without proving them in details, since there are exactly the same as in the aforementioned references.

Conditioned central limit theorem starting away from the boundary. Since S(n) has all moments uniformly bounded with respect to u, by [16], for all $r \ge 3, \gamma > 0$ there exists $\gamma > 0$ small enough and a coupling between $S_n - n\mu$ and a Brownian motion B^u with variance Σ^u such that

$$\mathbb{P}(\sup_{0 \leqslant t \leqslant n} |S_{\lfloor t \rfloor} - t\mu - B_t^u| > n^{1/2 - \gamma}) \leqslant C n^{-r},$$

with C independent of $u \in \Sigma$. As in [11, Lem. 20] we deduce from this by choosing $\gamma = 1/4 + 2\varepsilon$ and using Lemma 21 that for any $A \subset \mathbb{R}^2$ compact with non-empty interior, uniformly on $x \in \mathbb{H}_1$ with $x_2 \ge n^{1/4-\varepsilon}$ with ε small (only depending on γ, r),

(45)

$$\mathbb{P}_{u}(x+S(n) \in n\mu + \sqrt{n}A, \tau_{x}^{1} > n) \sim \int_{n\mu + \sqrt{n}A \cap \mathbb{R} \times \mathbb{R}_{\geq 0}} K_{n}^{B^{u}}(x,y) dy$$

$$\sim \frac{c(u)x_{2}\mu_{2}}{2\pi \Sigma_{11}^{u}\sqrt{\det(\Sigma^{u})}} \int_{A} \exp\left(-\frac{\|C^{u}y\|^{2}}{2}\right) dy,$$

(remark that we use the fact that $n\mu + \sqrt{n}A \in \mathbb{R} \times \mathbb{R}_{\geq 0}$ for n large enough, due to the condition $n^{-1/2} = o(u_2)$) and

(46)
$$\mathbb{P}_u(|S(n)| \ge t, \tau_x^1 > n) \le C x_2 \mu_2 \exp\left(-ct^2\right) dy + O(n^{-r}),$$

for some C, c > 0 and r large enough.

Conditioned central limit theorem starting close the boundary. Introduce the stopping times $\nu_n^x = \inf(m \ge 1, x_2 + S_2(m) \ge n^{1/4-\varepsilon})$. Then, as in [11, Lemma 14] $\mathbb{P}(\nu_n \ge n^{1/2-\varepsilon}) = O(e^{-Cn^{\varepsilon}})$ and by Lemma 13,

$$\mathbb{P}(\nu_n \leqslant n^{1/2-\varepsilon}, |S(\nu_n) - \nu_n \mu^u| \leqslant n^{1/4-\varepsilon/4}) \ge 1 - \exp(-Cn^{-\varepsilon'})$$

for some $\varepsilon' > 0$. Hence, we have by (45)

$$\begin{split} \mathbb{P}_{u}(x+S(n) \in n\mu + \sqrt{n}A, \tau_{x}^{1} > n) \\ = \mathbb{E} \left(\mathbb{P} \left(x + S(\nu_{n}^{x}) \in x + S(\nu_{n}^{x}) + (n - \nu_{n}^{x})\mu + \sqrt{n}(A + n^{-1/2}(\nu_{n}^{x}\mu - S(\nu_{n}^{x})), \tau_{n}^{x} < \tau_{x}^{1} \right) \\ \tau_{x+S(\nu_{n}^{x})}^{1} > n - \nu_{n}^{x} \right), \nu_{n}^{x} < \tau_{x}^{1} \right) \\ = \mathbb{E} \left(\mathbb{P} \left(x + S(\nu_{n}^{x}) \in x + S(\nu_{n}^{x}) + (n - \nu_{n}^{x})\mu + \sqrt{n}(A + o(n^{-1/2 - \frac{\varepsilon}{4}})), \tau_{x+S(\nu_{n}^{x})}^{1} > n - \nu_{n}^{x} \right), \nu_{n} < n^{1/2 - \varepsilon}, \nu_{n}^{x} < \tau_{x}^{1} \right) + o \left(\exp \left(-Cn^{-\varepsilon'} \right) \right) \\ \sim c(u) \mathbb{E} \left(x_{2} + S_{2}(\nu_{n}^{x}), \nu_{n} < n^{1 - \varepsilon}, \nu_{n}^{x} < \tau_{x}^{1} \right) \frac{\mu_{2}}{2\pi n \sqrt{\Sigma_{11}^{u} \det(\Sigma^{u})}} \int_{A} \exp \left(-\frac{\|C^{u}y\|^{2}}{2} \right) dy, \end{split}$$

uniformly on $u \in \Sigma$ and $x \in \mathbb{H}_1, x = o(\sqrt{n})$. Then, using that $\mathbb{P}(|x_2 + S_2(n)|, \nu_n \ge n^{1/2-\varepsilon}) = O(e^{-Cn^{\varepsilon'}})$ as in [11, Lem. 16] and the fact that $S_2(k) - k\mu_2^u$ is a martingale, $\nu_n^x \wedge \tau_x^1$ is a stopping time and $x_2 + S_2(\tau_x^1) = 0$ (because S is skip-free), we get

$$\mathbb{E}\left(x_2 + S_2(\nu_n^x), \nu_n < n^{1/2-\varepsilon}, \nu_n^x < \tau_x^1\right) = \mathbb{E}\left(x_2 + S_2(\nu_n^x \wedge n^{1/2-\varepsilon}), \nu_n^x < \tau_x^1\right) + O(e^{-Cn^{\varepsilon'}})$$
$$= \mathbb{E}\left(x_2 + S_2(\nu_n^x \wedge \tau_x^1 \wedge n^{1/2-\varepsilon})\right) + O(e^{-Cn^{\varepsilon'}})$$
$$= x_2 + \mu_2^u \mathbb{E}(\nu_n^x \wedge \tau_x^1 \wedge n^{1/2-\varepsilon}) + O(e^{-Cn^{\varepsilon}}).$$

Since $\mu_2^u \leq n^{-1/2+\eta}$, for η small enough we have

$$\mu_2^u \mathbb{E}(\nu_n^x \wedge \tau_x^1 \wedge n^{1/2-\varepsilon}) \leqslant n^{-1/2+\eta} n^{1/2-\varepsilon} = o(1),$$

uniformly on $u \in \Sigma$ in the range of the Lemma. Finally, uniformly on such u and $x \in \mathbb{H}_1$, $|x| = o(\sqrt{n})$,

$$\mathbb{P}_u(x+S(n)\in n\mu+\sqrt{n}A, \tau_x^1>n)\sim_{n\to\infty}\frac{c(u)x_2\mu_2}{2\pi n\Sigma_{11}^u\sqrt{\det(\Sigma^u)}}\int_A\exp\left(-\frac{\|C^uy\|^2}{2}\right)dy$$

Similarly, we get for $x \in \mathbb{H}_1$ and $t \ge 0$,

$$\mathbb{P}_u(|S(n) - n\mu^u| \ge \sqrt{n}t, \tau_x^1 > n) \le Cx_2\mu_2 \exp\left(-c^u t^2\right) dy + o(n^{-r}).$$

Local limit theorem starting close the boundary. Using the same proof as the one of [11, Thm 5] with the local limit theorem from Spitzer [27, Ch. 7 P10] which is valid for any random walk whose steps set generates \mathbb{Z}^2 , we deduce

$$\mathbb{P}_{u}(x+S(n)=y,\tau_{x}^{1}>n) = \frac{c(u)x_{2}\mu_{2}}{\Sigma_{11}^{u}2\pi n\sqrt{\det(\Sigma^{u})}}\exp\left(-\frac{\|C^{u}(y-n\mu^{u})\|^{2}}{2n}\right) + o\left(\frac{\mu_{2}^{u}x_{2}}{n}\right)$$

with $o(\cdot)$ uniform on u and $x, y \in \mathbb{H}_1$ with $|x| \leq \sqrt{n}$, and

$$\mathbb{P}_{u}(x+S(n)=y,\tau_{x}^{1}>n) \leqslant \frac{Cx_{2}\mu_{2}}{n} \exp\left(-c^{u}\|y-n\mu^{u}\|^{2}\right) dy + o\left(n^{-(1+r)}\right). \qquad \Box$$

Acknowledgments

Part of this work has been written during a stay at the University of Münster of the first and second authors, who would like to thank the Institute for Mathematical Stochastics for the very good working conditions. We thank the editors for their kind invitation to submit a paper to this special issue in honour of J. W. Cohen. We further thank Ivo Adan, Onno Boxma, Johan van Leeuwaarden and Michel Mandjes for continuous and fruitful interactions since 2010. We thank Ivo Adan for useful discussions at an early stage of the project. KR thanks Alin Bostan and Lucia Di Vizio for interesting discussions on singular random walks.

References

- I. J.-B. F. Adan (1991). A compensation approach for queueing problems. Dissertation, Technische Universiteit Eindhoven
- [2] I. J.-B. F. Adan, J. Wessels and W. H. M. Zijm (1990). Analysis of the symmetric shortest queue problem. *Comm. Statist. Stochastic Models* 6 691–713
- [3] I. J.-B. F. Adan, J. Wessels and W. H. M. Zijm (1993). A compensation approach for two-dimensional Markov processes. Adv. in Appl. Probab. 25 783–817
- [4] M. Bousquet-Mélou and M. Mishna (2010). Walks with small steps in the quarter plane. Algorithmic probability and combinatorics, 1–39, Contemp. Math., 520, Amer. Math. Soc., Providence, RI
- [5] M. Bousquet-Mélou and M. Petkovšek (2000). Linear recurrences with constant coefficients: the multivariate case. Formal power series and algebraic combinatorics (Toronto, ON, 1998). Discrete Math. 225 51–75
- [6] M. Bousquet-Mélou and M. Petkovšek (2003). Walks confined in a quadrant are not always D-finite. Theoret. Comput. Sci. 307 257–276
- [7] M. Brown, E. A. Pekoz and S. M. Ross (2010). Some results for skip-free random walk. Probab. Engrg. Inform. Sci. 24 491–507
- [8] J. W. Cohen (1992). Analysis of random walks. Studies in Probability, Optimization and Statistics, 2. IOS Press, Amsterdam
- [9] J. W. Cohen and O. J. Boxma (1983). Boundary value problems in queueing system analysis. North-Holland Mathematics Studies, 79. North-Holland Publishing Co., Amsterdam
- [10] M. Defosseux (2015). Affine Lie algebras and conditioned space-time Brownian motions in affine Weyl chambers. Probab. Theory Relat. Fields 165 649–665
- [11] D. Denisov and V. Wachtel (2015). Random walks in cones. Ann. Probab. 43 992-1044
- [12] V. Despax (2018). On some lattice random walks conditioned to stay in Weyl chambers. Markov Process. Related Fields 24 733–758
- [13] J. L. Doob (1959). Discrete potential theory and boundaries. J. Math, and Mech. 8 433-458
- [14] T. Dreyfus, C. Hardouin, J. Roques and M. F. Singer (2020). Walks in the quarter plane: genus zero case. J. Combin. Theory Ser. A 174 105251, 25 pp.
- [15] J. Duraj, K. Raschel, P. Tarrago and V. Wachtel (2022). Martin boundary of random walks in convex cones. Ann. Henri Lebesgue 5 559-609
- [16] U. Einmahl (1989). Extensions of results of Komlos, Major, and Tusnady to the multivariate case. J. Multivariate Anal. 28 20–68
- [17] G. Fayolle, R. Iasnogorodski and V. Malyshev (2017). Random walks in the quarter plane. Second edition. Probability Theory and Stochastic Modelling, 40. Springer, Cham
- [18] R. Garbit and K. Raschel (2014). On the exit time from a cone for Brownian motion with drift. *Electron. J. Probab.* 19, no. 63, 27 pp.
- [19] V. H. Hoang, K. Raschel and P. Tarrago (2022). Constructing discrete harmonic functions in wedges. Trans. Amer. Math. Soc. 375 4741–4782
- [20] I. Ignatiouk-Robert and C. Loree (2010). Martin boundary of a killed random walk on a quadrant. Ann. Probab. 38 1106–1142
- [21] I. Kurkova and K. Raschel (2011). Random walks in \mathbb{Z}^2_+ with non-zero drift absorbed at the axes. Bull. Soc. Math. France 139 341–387
- [22] C. Lecouvey and K. Raschel (2016). t-Martin boundary of killed random walks in the quadrant. Séminaire de Probabilités XLVIII, 305–323, Lecture Notes in Math., 2168, Springer, Cham
- [23] S. Melczer and M. Mishna (2014). Singularity analysis via the iterated kernel method. Combin. Probab. Comput. 23 861–888

- [24] M. Mishna and A. Rechnitzer (2009). Two non-holonomic lattice walks in the quarter plane. Theoret. Comput. Sci. 410 3616–3630
- [25] P. Ney and F. Spitzer (1966). The Martin boundary for random walk. Trans. Amer. Math. Soc. 121 116–132
- [26] K. Raschel and P. Tarrago (2020). Boundary behavior of random walks in cones. Markov Process. Related Fields 26 711–756
- [27] F. Spitzer (1976). Principles of random walks, 2nd Edition. Springer, Graduate Texts in Mathematics 34

Institut Denis Poisson, UMR CNRS 7013, Université de Tours et Université d'Orléans, Parc de Grandmont, 37200 Tours, France

Email address: viet.hung-hoang@lmpt.univ-tours.fr

LABORATOIRE ANGEVIN DE RECHERCHE EN MATHÉMATIQUES, UMR CNRS 6093, UNIVERSITÉ D'ANGERS, 2 BOULEVARD LAVOISIER, 49000 ANGERS, FRANCE

Email address: raschel@math.cnrs.fr

LABORATOIRE DE PROBABILITÉS, STATISTIQUE ET MODÉLISATION, UMR CNRS 8001, SORBONNE UNIVERSITÉ, 4 PLACE JUSSIEU, 75005 PARIS, FRANCE

Email address: pierre.tarrago@sorbonne-universite.fr